



HAL
open science

Gauss-Kronecker Curvature and equisingularity at infinity of definable families

Nicolas Dutertre, Vincent Grandjean

► **To cite this version:**

Nicolas Dutertre, Vincent Grandjean. Gauss-Kronecker Curvature and equisingularity at infinity of definable families. 2019. hal-02308350

HAL Id: hal-02308350

<https://univ-angers.hal.science/hal-02308350v1>

Preprint submitted on 25 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

GAUSS-KRONECKER CURVATURE AND EQUISINGULARITY AT INFINITY OF DEFINABLE FAMILIES

NICOLAS DUTERTRE AND VINCENT GRANDJEAN

ABSTRACT. Assume given a polynomially bounded o-minimal structure expanding the real numbers. Let $(T_s)_{s \in \mathbb{R}}$ be a globally definable family of C^2 -hypersurfaces of \mathbb{R}^n . Upon defining the notion of generalized critical value for such a family, we show that the functions $s \rightarrow |K(s)|$ and $s \rightarrow K(s)$, respectively the total absolute Gauss-Kronecker and total Gauss-Kronecker curvature of T_s , are continuous in any neighbourhood of any value which is not generalized critical. In particular this provides a necessary criterion of equisingularity for the family of the levels of a real polynomial.

1. INTRODUCTION

One of the main goal of equisingularity theory (of families of subsets, functions, mappings) is to find relations between numerical data and regularity conditions. In the local complex analytic case, this subject has been widely studied since the end of the 60's and many interesting results, some of them now classical, have been established. For example, Hironaka [Hir] proved that the multiplicity is constant along the strata of a Whitney stratification of a complex analytic set. In [Tei1] Teissier proved that a μ^* -constant family of hypersurfaces with isolated singularities is Whitney equisingular. The reverse implication was proved later by Briançon and Speder [BrSp]. These results were extended to the case of ICIS by Gaffney [Gaf1]. Maybe the most important result of local complex analytic equisingularity theory is Teissier's polar equimultiplicity theorem [Tei2], which states that Whitney regularity is equivalent to constancy of polar multiplicities. Teissier's results were refined and extended by Gaffney [Gaf2] to obtain sufficient conditions for equisingularity of a family of mappings.

When one considers global equisingularity problems, the first natural family to study is the family of fibres of a polynomial mapping. Following [Tho], a polynomial function from \mathbb{K}^n to \mathbb{K} , for $\mathbb{K} = \mathbb{R}$ or \mathbb{C} , is a smooth locally trivial fibration above the connected components of the complement of a (minimal) finite subset $B(f)$ of \mathbb{K} , called the set of bifurcation values of f . In the complex plane case, Hà and Lê [HL] gave the following numerical criterion to characterize bifurcation values: *A value c does not lie in $B(f)$ if and only if the Euler characteristic of the fibres of f is constant in a neighborhood of c .* This result was generalized by Parusiński [Par] to the case of complex polynomials with isolated singularities at infinity, and then by Siersma and Tibăr [SiTi1] to the case of complex polynomials with isolated \mathcal{W} -singularities at infinity. In [Tib1] Tibăr studies the more general situation of a 1-parameter family of complex hypersurfaces, and proves a global version of the results of Teissier and Briançon and Speder mentioned above: Considering a family of complex affine hypersurfaces $X_\tau = \{x \in \mathbb{C}^n : F(\tau, x) = 0\}$ given by a polynomial function $F : \mathbb{C} \times \mathbb{C}^n \mapsto \mathbb{C}$, he defines the notion of t -equisingularity at infinity and proves, under some additional conditions, that t -equisingularity at infinity is controlled by the constancy of a finite sequence of numbers, called the generic polar intersection multiplicities. As a consequence, if the family consists of non-singular affine hypersurfaces, then the constancy of the generic polar intersection multiplicities at τ_0 implies that the family is C^∞ trivial at τ_0 .

Date: February 26, 2019.

2010 Mathematics Subject Classification. Primary 14P10, Secondary 57R70 03C64.

Key words and phrases. Gauss-Kronecker curvature, total curvatures, generalized critical values, definable families.

Both authors were partially supported by the ANR project LISA 17-CE400023-01

Vincent Grandjean was supported by CNPq-Brazil grant 150555/2011-3 and FUNCAP/CAPES/CNPq-Brazil grant 305614/2015-0.

In the real semi-algebraic/sub-analytic setting (or more generally in the definable setting), it is hopeless to expect that *constancy of numerical data is equivalent to regularity conditions*. First, because of lack of connectivity, one cannot define invariants like the μ^* -sequence, polar multiplicities or generic polar intersection multiplicities. However, using arguments from differential topology and integral geometry, one sees that these invariants admit geometric characterizations that still make sense in the real case. For instance, the multiplicity of a complex analytic germ is equal to its density [Dra] and the μ^* -sequence, the polar multiplicities and the generic polar intersection multiplicities are related to curvature integrals (see [La, Loe, Dut1, SiTi2]). Unfortunately, in the real situation, these geometric quantities do not belong to discrete sets and therefore, one cannot expect results relating their constancy to regularity conditions. It is more reasonable to study properties like continuity or Lipschitz continuity in the parameters of the family. The first result in this direction is due to Comte [Com], who established a real version of Hironaka's theorem, proving that the density is continuous along the strata of a (w) -stratification of a sub-analytic set. This result was generalized and strengthened by Valette [Val]: continuity of the density holds for (b) -regular stratifications and the density is Lipschitz continuous along the strata of (w) -stratifications. Later Comte and Merle [ComMe] established a real version of Teissier's theorem [Tei2]. Using tools from integral geometry and geometric measure theory, they associated with each sub-analytic germ a sequence of numbers, called the local Lipschitz-Killing invariants, and showed that they are continuous along the strata of a (w) -stratification of a sub-analytic set. Recently, Nguyen and Valette [NgVa] extended this continuity result to (b) -stratifications and moreover proved that these invariants are Lipschitz continuous along the strata of a (w) -stratification (see also the first author work [Dut2] for relations with the densities of polar images).

In the global real context, it is still true that the bifurcation set of a definable function from \mathbb{R}^n to \mathbb{R} is a finite set of points (see [NeZa, LoZa, Tib2, d'Ac1]). In [TiZa] Tibăr and Zaharia provided necessary and sufficient conditions for a real plane polynomial function to be locally trivial over the neighborhood of a regular value (see [JoTi] for a generalization to a family of real curves). Unlike the complex case, their criterion is not only numerical but involves topological conditions at infinity. Later in [CosPe], Coste and de la Puente proved an equivalent version of Tibăr-Zaharia's results in terms of polar curves. Due to the links between polar curves and the Gauss-Kronecker curvature of the levels of a function provided by exchange formulas, it seems natural to study the variations of the total curvature of the levels (i.e. the integral of the Gauss-Kronecker curvature on the level) of a definable function, and to seek how bifurcation values interfere in these variations.

That is what the second author did in two papers. In [Gra1] he considers a globally definable function $f : \mathbb{R}^n \mapsto \mathbb{R}$ of class at least C^2 , and proved that the following functions:

$$t \mapsto \int_{f^{-1}(t)} \kappa, \quad \text{and} \quad t \mapsto \int_{f^{-1}(t)} |\kappa|$$

where κ is the Gauss-Kronecker curvature, admit at most finitely many discontinuities. In [Gra2] he proved that if the function $t \mapsto \int_{f^{-1}(t)} |\kappa|$ is continuous at a regular value c which satisfies an extra condition, then c is not a bifurcation value of f . He explained that for a real polynomial function with isolated singularities at infinity this extra condition is always satisfied, so this result can be interpreted as a real version of Parusiński's result mentioned above.

The aim of the present paper is to provide a kind of reverse implication of the latter mentioned result.

We will work in the more general situation of a one parameter family of hypersurfaces. More precisely, we consider a globally definable function over an a priori given polynomially bounded o -minimal structure $F : \mathbb{R}^n \times \mathbb{R} \mapsto \mathbb{R}$ of class C^{2+m} with non-negative integer m . Assuming that 0 is a regular value of F , the 0-level $M = F^{-1}(0)$ is thus a globally definable hypersurface in \mathbb{R}^{n+1} of class C^{2+m} . We use the coordinates (\mathbf{x}, t) in $\mathbb{R}^n \times \mathbb{R}$ and we write $\mathbf{t}_M : M \mapsto \mathbb{R}$, $(\mathbf{x}, t) \mapsto t$ for the projection on the t -axis.

For a value c in \mathbb{R} , let $M_c = \mathbf{t}_M^{-1}(c)$ and $T_c = \pi_M(M_c) \subset \mathbb{R}^n$, where π_M is the projection from M to \mathbb{R}^n . If c is a regular value, then the hypersurface T_c is oriented by $\partial_{\mathbf{x}}F(\mathbf{x}, c)$. Therefore, we consider the

Gauss-Kronecker curvature κ_c of T_c and define two functions:

$$c \mapsto K(c) = \int_{T_c} \kappa_c(\mathbf{p}) d\mathbf{p}, \quad \text{and} \quad c \mapsto |K|(c) = \int_{T_c} |\kappa_c|(\mathbf{p}) d\mathbf{p}.$$

By a straightforward adaptation of the methods of [Gra1], we show that these two functions have finitely many discontinuities (Theorem 6.1) and in Theorem 8.1, we give a criterion on the regular value c of \mathbf{t}_M for the function $t \mapsto |K|(t)$ to be continuous at c . Namely, we prove

Theorem 8.1. *Let c be a regular value taken by \mathbf{t}_M at which it is horizontally spherical at infinity. Then the total absolute curvature function $t \mapsto |K|(t)$ is continuous at c . Consequently the total curvature function $t \mapsto K(t)$ is continuous at c .*

The notion of *horizontally spherical at infinity* is a regularity condition at infinity: A regular value c of \mathbf{t}_M is horizontally spherical at infinity if for any sequence $(\mathbf{p}_k)_{k \in \mathbb{N}}$ of M converging at infinity to (\mathbf{u}, c) , \mathbf{u} is orthogonal to the limit of the unitary gradients $\frac{\nabla \mathbf{t}_M}{|\nabla \mathbf{t}_M|}(\mathbf{p}_k)$. A key ingredient of the proof of our main result, Theorem 8.1 is Lemma 8.2 stating, informally, that *under these hypotheses there no accumulation of curvature at infinity nearby the level c .*

We also prove that (t) -equisingularity at infinity implies horizontal sphericalness (Corollary 5.2). Therefore Theorem 8.1 shows that (t) -equisingularity at infinity implies continuity of the function $t \mapsto |K|(t)$. This can be considered as a first step towards a real version of Tibř's result [Tib1] mentioned above.

To be complete, we show here more than Theorem 8.1. Its conclusion also holds true in any connected component of the pencil of levels over a small interval of regular values $]c - \varepsilon, c + \varepsilon[$ (see Theorem 8.3). In other words the connected components of the pencil of levels cannot compensate altogether the a priori possible discontinuities of some.

The paper is organized as follows. Section 2 contains material on compactifications, o -minimal structures and Thom's (a_f) condition. In Section 3, we recall some facts about conormal geometry so that we can introduce the notion of t -equisingularity. Sections 4, 5 and 7 contain definitions and new results on regularity at infinity of globally definable C^{2+m} -families of hypersurfaces. In Section 6, we generalize the results of [Gra1] to our situation. Section 8 contains the proof of the main result. Section 9 deals with the particular case of the levels of a function.

Acknowledgments. The authors are very grateful to Si Tiep Dinh for useful, fruitful and inspiring conversations. The second author would like to thank the I2M and LAREMA for their working conditions while visiting the first author.

2. MISCELLANEOUS MATERIAL

Let \mathbb{R}^n be the Euclidean space of positive dimension n .

Let $\langle -, - \rangle$ be the associated scalar product. For any point \mathbf{x} of \mathbb{R}^n , let $|\mathbf{x}|$ be the norm $\sqrt{\langle \mathbf{x}, \mathbf{x} \rangle}$ of \mathbf{x} .

Let \mathbf{S}_R^{q-1} be the Euclidean sphere of \mathbb{R}^q of center the origin and positive radius R .

Let \mathbf{B}_R^q be the closed Euclidean ball of \mathbb{R}^q of center the origin and positive radius R . When q is understood we will only write \mathbf{B}_R .

Let $\text{clos}(-)$ denote the operation "taking the closure of" in \mathbb{R}^n . Each Euclidean space \mathbb{R}^n embeds semi-algebraically in the closed unit-ball \mathbf{B}_1^n , as its interior via the mapping $\mathbf{x} \mapsto \frac{\mathbf{x}}{\sqrt{1+|\mathbf{x}|^2}}$. We may then speak of \mathbf{B}_1^n as the *spherical compactification* of \mathbb{R}^n (see the next section).

Let \mathcal{M} be an o -minimal structure expanding the real field \mathbb{R} . Assume it is polynomially bounded and let $\mathbb{F}_{\mathcal{M}}$ be the field of its exponents ([vdDM, vdD]). Any subset of any \mathbb{R}^p definable in \mathcal{M} will be called below *definable*.

Usually globally definable subsets of \mathcal{M} are defined as definable subsets of any closed unit Euclidean ball.

Since each \mathbb{R}^n embeds semi-algebraically in the closed unit-ball \mathbf{B}_1^n , a subset X of \mathbb{R}^n is *globally definable* if it is definable in the spherical compactification of \mathbb{R}^n .

Let X be a subset of \mathbb{R}^n . A mapping $f : X \mapsto \mathbb{R}^p$ is globally definable if its graph is globally definable in \mathbb{R}^{n+p} .

We would like to remind the following fact (see [d'Ac1]): *Let $\gamma : [1, +\infty[\mapsto \mathbb{R}^n$ be a C^1 globally definable arc such that $\gamma(t) \mapsto \infty$ as t goes to $+\infty$. Then there exists a unit vector \mathbf{u} of \mathbf{S}^{n-1} such that*

$$\lim_{\infty} \frac{\gamma}{|\gamma|} = \mathbf{u} = \lim_{\infty} \frac{\gamma'}{|\gamma'|}.$$

Let $f : (\mathbb{R}_{\geq 1}, +\infty) \rightarrow \mathbb{R}$ be the germ at ∞ of a continuous globally definable function. We write $f \sim t^e$ for an exponent e in $\mathbb{F}_{\mathcal{M}} \cup \{-\infty\}$, with the convention that $t^{-\infty} = 0$ for large t , to mean

$$f \sim t^e \iff \lim_{t \rightarrow +\infty} \frac{f(t)}{t^e} \in \mathbb{R}^*.$$

Note that there always exists such an exponent e .

Let $\mathbf{G}(p, n)$ be the Grassmann manifold of p -vector subspaces of \mathbb{R}^n . We denote $\mathbf{G}^\vee(p, n)$ the space of p -vector subspaces of the space $L(\mathbb{R}^n, \mathbb{R})$ of linear forms over \mathbb{R}^n , and we will call it sometimes the *dual* of $\mathbf{G}(n-p, n)$.

We recall *Thom's condition* (or relative Whitney's condition (a)).

Let X, Y be two connected C^1 submanifolds of a definable compactification of \mathbb{R}^n , such that Y is contained in $\text{clos}(X) \setminus X$. Let $g : (X \sqcup Y) \mapsto \mathbb{R}$ be a C^1 mapping, for $X \sqcup Y$ the disjoint union of X and Y . Let \mathbf{y} be a point of Y .

The function g satisfies *Thom (a_g)-condition at \mathbf{y}* if the following two conditions hold:

- (i) For any sequence $(\mathbf{x}_k)_k$ of points of X converging to \mathbf{y} such that the sequence $(T_{\mathbf{x}_k} X)_k$ converges to T in the appropriate Grassmann bundle, then $T_{\mathbf{y}} Y$ is contained in T ;
- (ii) For any sequence $(\mathbf{x}_k)_k$ of points of X converging to \mathbf{y} , such that the sequence $(T_{\mathbf{x}_k} X)_k$ converges to T which contains $T_{\mathbf{y}} Y$ and the sequence $(\ker d_{\mathbf{x}_k} g)_k$ converges to K in the appropriate Grassmann bundle, then $\ker d_{\mathbf{y}} g$ is contained in K .

In practice we want to stratify g with Thom's condition asking that the strata Y is contained in some specified level of g .

3. COMPACTIFICATION AND CONORMAL GEOMETRY AND t -EQUISINGULARITY AT INFINITY

Let $\mathbf{0}$ be the origin of \mathbb{R}^n .

As already seen in the previous section, we can compactify \mathbb{R}^n as the closed unit ball \mathbf{B}_1^n .

An alternative presentation to the spherical compactification is the spherical blowing-up \mathbf{bl}_∞ of \mathbb{R}^n at infinity, that is the mapping given by

$$\begin{aligned} \mathbf{bl}_\infty : \mathbf{S}^{n-1} \times]0, +\infty[&\mapsto \mathbb{R}^n \setminus \{\mathbf{0}\} \\ (\mathbf{u}, r) &\mapsto \frac{\mathbf{u}}{r} \end{aligned}$$

It is a Nash diffeomorphism and a re-parametrization of $\mathbb{R}^n \setminus \{\mathbf{0}\}$ embedded in \mathbf{B}_1^n . It is more convenient to look at it this way since it is a good real avatar of the projective compactification $\mathbf{P}\mathbb{R}^n$ (which in our globally definable context is not as relevant as in the algebraic case).

We denote by $\mathbf{S}_\infty^{n-1} := \mathbf{S}^{n-1} \times 0$ the sphere at infinity. Let us denote and identify

$$\overline{\mathbb{R}^n} := \mathbb{R}^n \sqcup \mathbf{S}_\infty^{n-1} = (\mathbf{S}^{n-1} \times [0, +\infty]) \sqcup \mathbf{0},$$

the spherical compactification of \mathbb{R}^n at infinity, with boundary $\partial \overline{\mathbb{R}^n} := \mathbf{S}_\infty^{n-1}$, the sphere at infinity.

Let \overline{Z} be the closure of the subset Z of \mathbb{R}^n taken into $\overline{\mathbb{R}^n}$. The *tangent link of Z at ∞* is defined as

$$Z^\infty := \overline{Z} \cap \mathbf{S}_\infty^{n-1}.$$

The *tangent cone of Z at infinity* $C_\infty(Z)$ is defined as the (non-negative) cone over Z^∞ . Whence Z^∞ is not empty (equivalently Z is not bounded) we also observe that

$$Z^\infty := \text{clos} \left\{ \mathbf{u} \in \mathbf{S}^{n-1} : \exists Z \ni (x_k)_k \rightarrow \infty \text{ such that } \frac{x_k}{|x_k|} \rightarrow \mathbf{u} \right\}.$$

For any definable subset Z of \mathbb{R}^n , it is globally definable if, by definition, \overline{Z} is definable in $\overline{\mathbb{R}^n}$. Thus whenever a subset Z of \mathbb{R}^n is globally definable, its *tangent link at infinity* Z^∞ is definable and of dimension at most $\dim Z - 1$.

Although heavy to define it is convenient to use the formalism of conormal geometry. We are especially interested in conormal geometry at infinity.

Now let $Z = \{(\mathbf{x}, t) \in \mathbb{R}^n \times \mathbb{R} : G(\mathbf{x}, t) = 0\}$, where $G : \mathbb{R}^n \times \mathbb{R} \mapsto \mathbb{R}$ is a globally definable function of class at least C^2 , and let \overline{Z} be its closure in $\overline{\mathbb{R}^n} \times \mathbb{R}$.

We assume that 0 is a regular value of G and we consider Z as a definable family $\{Z_t\}_{t \in \mathbb{R}}$ of hypersurfaces in \mathbb{R}^n . Let $g : Z \rightarrow \mathbb{R}$ be a definable function which we assume to be C^1 . For any regular point (\mathbf{x}, t) of the function g , let $T_{(\mathbf{x}, t)}g$ be the subspace of $T_{(\mathbf{x}, t)}Z$ tangent at (\mathbf{x}, t) to the level of g through (\mathbf{x}, t) . Let us define the following subset of $\overline{\mathbb{R}^n} \times \mathbb{R} \times \mathbf{G}^\vee(1, n+1)$:

$$\mathcal{X}_g^\vee := \text{clos}\{(\mathbf{x}, t, \xi) \in Z \setminus \text{crit}(g) \times \mathbf{G}^\vee(1, n+1) : \xi(T_{(\mathbf{x}, t)}g) = 0\},$$

where $\mathbf{G}^\vee(1, n+1)$ is the dual of $\mathbf{G}(1, n+1)$.

Definition 3.1. *The relative conormal space of g is the space \mathcal{X}_g^\vee .*

Let $\pi_{n+1} : \overline{\mathbb{R}^n} \times \mathbb{R} \times \mathbf{G}^\vee(1, n+1) \mapsto \overline{\mathbb{R}^n} \times \mathbb{R}$ be the projection given as $\pi_{n+1}(\mathbf{x}, t, \xi) = (\mathbf{x}, t)$.

Definition 3.2. *The relative conormal space of g at infinity is the space \mathcal{X}_g^∞ defined as*

$$\mathcal{X}_g^\infty := \pi_{n+1}^{-1}(Z^\infty) \cap \mathcal{X}_g^\vee,$$

where $Z^\infty = \overline{Z} \cap (\mathbf{S}_\infty^{n-1} \times \mathbb{R})$.

For any $\mathbf{p} \in Z^\infty$, let $(\mathcal{X}_g^\vee)_\mathbf{p}$ be the fibre of \mathcal{X}_g^\vee above \mathbf{p} , that is $(\mathcal{X}_g^\vee)_\mathbf{p} = \pi_{n+1}^{-1}(\mathbf{p}) \cap \mathcal{X}_g^\vee$.

We introduce now the notion of t -equisingularity [Tib1] adapted to the context of Section 4.

Let $\mathbf{r}_Z : Z \mapsto \mathbb{R}$ be defined as $(\mathbf{x}, t) \mapsto |\mathbf{x}|$. It is continuous globally definable and C^1 outside $0 \times \mathbb{R} \cap Z$.

The *space of characteristic covectors \mathcal{C} of Z at infinity* is the subset of $\overline{\mathbb{R}^n} \times \mathbb{R} \times \mathbf{G}^\vee(1, n+1)$ defined as

$$\mathcal{C}(Z) := \mathcal{X}_{\mathbf{r}_Z}^\infty.$$

It is closed and definable.

Let $\tau : \overline{\mathbb{R}^n} \times \mathbb{R}$ be defined as $(\mathbf{x}, t) \mapsto t$.

The following notion is due to Tibăr [Tib1]:

Definition 3.3. *Let \mathbf{p} in Z^∞ .*

(i) *The family $\{Z_t\}_{t \in \mathbb{R}}$ is t -equisingular at \mathbf{p} if*

$$\mathcal{C}(Z)_\mathbf{p} \cap (\mathcal{X}_\tau^\infty)_\mathbf{p} = \emptyset.$$

(ii) *The family $\{Z_t\}_{t \in \mathbb{R}}$ is t -equisingular at infinity at c if it is t -equisingular at \mathbf{p} for all \mathbf{p} in $Z^\infty \cap \tau^{-1}(c)$.*

The definition above is slightly different from those given in [SiTi1, Tib1, DiRuTi], since there it is given via the projective compactification of \mathbb{R}^n . Anyhow they are equivalent.

Any co-vector in $(\mathcal{X}_\tau^\infty)_\mathbf{p}$ has kernel the horizontal hyperplane $\mathbb{R}^n \times 0$. We deduce that any limit of tangent spaces T of \mathbf{r}_Z at \mathbf{p} does not lie in $\mathbb{R}^n \times 0$ whenever $\{Z_t\}_{t \in \mathbb{R}}$ is t -equisingular at \mathbf{p} .

4. REGULARITIES AT INFINITY FOR DEFINABLE FAMILIES OF HYPERSURFACES

We present here two regularity conditions at infinity for the function restriction of a coordinate projection along a globally definable one parameter family of hypersurfaces. In the next section, we will compare altogether these regularity conditions with t -equisingularity, introduced in the previous section.

Let $F : \mathbb{R}_x^n \times \mathbb{R}_t \mapsto \mathbb{R}$ be a C^{2+m} globally definable function, for some non-negative integer m .

Assuming that $\mathbb{R}^n \times \mathbb{R}$ is equipped with the canonical Euclidean structure, let ∇F be the gradient field of F . Without further hypotheses, the real number 0 may be a critical value of F , and ∇F may be vanishing on the zero level of F .

Working Hypothesis: 0 is a regular value of F .

Let M be the zero locus $F^{-1}(0)$ of the function F , which is a closed globally definable subset of \mathbb{R}^{n+1} and a C^{2+m} hypersurface. Let \overline{M} be its closure in $\overline{\mathbb{R}^n} \times \mathbb{R}$.

We define two mappings $(\pi, \mathbf{t}) : \overline{\mathbb{R}^n} \times \mathbb{R} \mapsto \overline{\mathbb{R}^n} \times \mathbb{R}$ obtained respectively as the restrictions of the projection over $\overline{\mathbb{R}^n}$ and over \mathbb{R} , and both are semi-algebraic.

Let \mathbf{t}_M be $\mathbf{t}|_M$ the restriction of \mathbf{t} to M and let π_M be the restriction of π to M , both are C^{2+m} and globally definable mappings. Let us write $M_c := \mathbf{t}_M^{-1}(c)$ and $T_c := \pi_M(M_c)$ subset of \mathbb{R}^n .

Definition 4.1. Let c be a value taken by \mathbf{t}_M . The function \mathbf{t}_M is said locally C^k trivial at c if there exists a positive real number ε such that $\mathbf{t}_M^{-1}(]c - \varepsilon, c + \varepsilon[)$ is a trivial C^k -bundle with fibre M_c .

Mimicking what was done for level hypersurfaces of functions [LoZa, TiZa, d'Ac1, d'AcGr1, d'AcGr2, Gra2], sufficient conditions about the gradient of \mathbf{t}_M guarantee trivialization (see below). Since M is globally definable and each of its connected component is orientable, let ν_M be a C^{1+m} globally definable unitary field normal to M . Since 0 is not a critical value of F , we choose

$$\nu_M := \frac{\nabla F}{|\nabla F|} = \nu_M^x + \nu_M^t \partial_t$$

where ν_M^x is the component of ν_M in $\mathbb{R}^n \times 0$, and writing $\nabla F = \partial_x F + \partial_t F \partial_t$, where $\partial_x F$ lies in $\mathbb{R}^n \times 0$.

Let $\mathbf{p} = (\mathbf{x}, t)$ be a point of M . We have

$$T_{\mathbf{p}}M = \{(\mathbf{u}, w) \in \mathbb{R}^n \times \mathbb{R} : \langle \partial_x F, \mathbf{u} \rangle + \partial_t F \cdot w = 0\}.$$

It is easy to prove the following relation:

$$\nabla \mathbf{t}_M = -\frac{\partial_t F}{|\nabla F|^2} \partial_x F + \frac{|\partial_x F|^2}{|\nabla F|^2} \partial_t = -\nu_M^t \nu_M^x + |\nu_M^x|^2 \partial_t$$

and thus

$$|\nabla \mathbf{t}_M| = \frac{|\partial_x F|}{|\nabla F|} = |\nu_M^x|.$$

The critical locus of \mathbf{t}_M is

$$\text{crit}(\mathbf{t}_M) = \{\mathbf{p} \in M : \partial_x F(\mathbf{p}) = 0\}.$$

Since M is a C^{2+m} orientable hypersurface, the function \mathbf{t}_M is C^{2+m} as well. Since it is globally definable, the set of its critical values $K_0(\mathbf{t}_M) := \mathbf{t}_M(\text{crit}(\mathbf{t}_M))$ is finite.

Let $\nu_{\mathbf{t}_M} : M \setminus \text{crit}(\mathbf{t}_M) \mapsto \mathbf{S}^n$ be the unitary gradient of $\nabla \mathbf{t}_M$,

$$\nu_{\mathbf{t}_M} := \frac{\nabla \mathbf{t}_M}{|\nabla \mathbf{t}_M|}.$$

The Local Conical Structure Theorem ensures the existence of a positive number S_M such that for any $S > S_M$ the hypersurface M is transverse with \mathbf{S}_S^n , the Euclidean sphere of radius S . As a consequence of this fact we also have:

Lemma 4.2. *For any $A > \max_{c \in K_0(\mathbf{t}_M)} |c|$, there exists R_A such that for any $R > R_A$ the globally definable C^{2+m} hypersurface $M \cap (\mathbb{R}^n \times]-A, A[)$ is transverse to the cylinder $\mathbf{S}_R^{n-1} \times \mathbb{R}$.*

Proof. Let $A \gg 1$ be given. Let us define the following subset

$$\Sigma := \{(\mathbf{x}, t) \in M : \mathbf{x} \wedge \nabla F(\mathbf{x}, t) = 0\}.$$

Note that $\Sigma \setminus \mathbf{0} \times \mathbb{R}$ is contained in $M \cap \{\partial_t F = 0\}$ and that Σ is a closed globally definable subset of M .

Let us assume that the statement of the lemma is not true. Thus there exists a C^1 globally definable path $\gamma : [1, +\infty[\rightarrow \Sigma \cap \mathbb{R}^n \times]-A, A[$ such that $\gamma(s) \rightarrow (\mathbf{u}, c)$ in M^∞ as s goes to $+\infty$, with $|c| < A$.

We can parameterize γ in such a way that $|\gamma(s)| = s$, which gives the following

$$\gamma(s) = (s\mathbf{u}, 0) + s^e \mathbf{v}(s)$$

for a C^1 and globally definable mapping $s \mapsto s^e \mathbf{v}(s) \in \mathbb{R}^n \times \mathbb{R}$ such that $\lim_{s \rightarrow \infty} \mathbf{v} \neq (0, 0)$ and $e < 1$. We also have that $s \mapsto \nu_M(s) = \frac{\nabla F}{|\nabla F|}(\gamma(s))$ goes to ν in \mathbf{S}^n as s goes to ∞ . Note that $\frac{\gamma'(s)}{|\gamma'(s)|}$ goes to $(\mathbf{u}, 0)$ as s goes to $+\infty$. Since

$$\gamma(s) \wedge \nu_M(s) = 0 \quad \text{and} \quad \langle \gamma'(s), \nu_M(s) \rangle = 0,$$

we deduce that

$$\mathbf{u} \wedge \nu = 0 \quad \text{and} \quad \langle \mathbf{u}, \nu \rangle = 0,$$

which is absurd. □

We can introduce now the Malgrange regularity condition at infinity.

Definition 4.3. *Let $c \in \mathbb{R}$ be a value.*

(i) *The function \mathbf{t}_M satisfies the Malgrange condition at c if there exist positive constants R, ε, A_c such that*

$$(4.1) \quad |\mathbf{x}| > R, |t - c| < \varepsilon \implies |\mathbf{x}| \cdot |\nabla \mathbf{t}_M(\mathbf{x}, t)| \geq A_c$$

which is equivalent to

$$(4.2) \quad |\mathbf{x}| > R, |t - c| < \varepsilon \implies |\mathbf{x}| \cdot |\partial_{\mathbf{x}} F| \geq A_c |\nabla F|.$$

(ii) *A value c which is not satisfying the Malgrange condition is called an asymptotic critical value (ACV for short). Let $K_\infty(\mathbf{t}_M)$ be the set of ACV of \mathbf{t}_M .*

Similarly to the case of real or complex polynomial families [Par, Tib1, Tib2, TiZa] we find

Theorem 4.4 (see also [LoZa, Kur, d'Ac1, d'AcGr1]). (i) *There exists a finite subset $B(\mathbf{t}_M)$ of \mathbb{R} such that the function \mathbf{t}_M is a locally C^{1+m} trivial at any value c not lying in $B(\mathbf{t}_M)$.*

(ii) $B(\mathbf{t}_M) \subset K_0(\mathbf{t}_M) \cup K_\infty(\mathbf{t}_M)$.

(iii) $K_\infty(\mathbf{t}_M)$ is finite.

(iv) *If c is a regular value taken by \mathbf{t}_M and does not lie in $K_\infty(\mathbf{t}_M)$, the local trivialization can be realized by a vector field colinear to $\nu_{\mathbf{t}_M}$.*

Proof. We are going to sketch the proofs of (iii) following [d'Ac1] and (iv) following [d'Ac1, d'Ac2, d'AcGr1]. Both (i) and (ii) can be deduced from these two points.

For simplicity we write f for \mathbf{t}_M .

Since \mathcal{M} is polynomially bounded, there exists a globally definable function $\rho : [1, +\infty[\rightarrow \mathbb{R}_+$ such that (see [d'Ac1, Lemma 3.3]):

$$(1) \lim_{R \rightarrow \infty} R^{-1} \rho(R) = +\infty \quad \text{and} \quad (2) K_\infty(f) = K_\infty^\rho(f)$$

where

$$K_\infty^\rho(f) := \{c \in \mathbb{R} : \exists (\mathbf{x}_k, t_k) \in M, \mathbf{x}_k \rightarrow \infty \text{ and } t_k \rightarrow c \text{ such that } \rho(|\mathbf{x}_k|) \cdot |\nabla f(\mathbf{x}_k, t_k)| \rightarrow 0\}.$$

In particular for R large enough there exists an exponent α of $(\mathbb{F}_M)_{>1}$ such that $\rho(R) \geq R^\alpha$.

Following the steps of [d'Ac1, Theorem 3.4], we show that $K_\infty^\rho(f)$ is finite.

Assume that there exists $c' > c$ such that for each $c \leq t \leq c'$ the level $\{f = t\}$ is neither empty nor is a critical level. Let us consider the following subset

$$\Delta := \{(s, w) \in \mathbb{R}^2 : \forall \delta > 0, \exists \varepsilon > 0, \exists R > 0 : |\mathbf{x}| > R, |\mathbf{t}_M(x) - w| < \varepsilon \implies |\mathbf{x}| \cdot |\nabla \mathbf{t}_M(\mathbf{x})| - s| < \delta\}.$$

This subset is globally definable. Let $\theta : [c, c'] \mapsto [0, +\infty[$ be the function defined as follows

$$\theta(t) := \inf\{\Delta \cap \{w = t\}\}.$$

It is globally definable. We wish to show that it vanishes only finitely many times on $[c, c']$. Assume that θ is identically 0 over $[c, c']$ (up to work with a smaller c'). Under these hypotheses the globally definable subset

$$\Sigma := \{\rho(|\mathbf{x}|) \cdot |\nabla f(\mathbf{x}, t)| < f(\mathbf{x}, t) - c\}$$

is not empty outside of a compact subset of $M \cap \mathbf{t}_M^{-1}[c - \varepsilon, c + \varepsilon]$ (see [d'Ac1, p. 40]). By definition of Σ , there exists a C^1 globally definable arc going to infinity $\gamma : [1, +\infty[\mapsto \Sigma$ such that

$$\lim_{\infty} f \circ \gamma = c.$$

Let $h = f \circ \gamma$, and let us parameterize γ such that $|\mathbf{x}(\gamma(R))| = R$, so that $|\gamma'|$ goes to 1 at infinity. We find

$$0 < -h' \leq \frac{h}{|\rho|} |\gamma'| < 2 \frac{h}{|\rho|}.$$

Let R_0 be large enough and let $u(R) = 2h(R_0) - h(R)$ once $R \geq R_0$, and let $a > 1$ be such that

$$\lim_{\infty} R^{-a} \rho(R) = +\infty.$$

We deduce for $R \geq R_0$ (up to taking a larger R_0)

$$0 < u'(R) < 2 \frac{u(R)}{|\rho(R)|} < \frac{u}{R^a}.$$

Applying the Gronwall Lemma provides for $R \geq R_0$

$$u(R) \leq h(R_0) \cdot \lambda(R_0) \text{ with } \lambda(R_0) := \exp\left(\frac{1}{(a-1)R_0^{a-1}}\right).$$

We know that $\lim_{\infty} u = 2h(R_0)$ but we can choose R_0 a priori such that $\lambda(R_0) < 2$, concluding that the function θ cannot vanish identically over $[c, c']$.

Point (iv) is of importance for the rest of the paper so we sketch its proof as a variation of the proof of [d'AcGr1, Theorem 3.5]. Let

$$\chi := \frac{1}{|\nabla \mathbf{t}_M|^2} \nabla \mathbf{t}_M.$$

Any trajectory γ of χ is parameterized by the levels of \mathbf{t}_M : starting at a point of M_c we find

$$\mathbf{t}_M(\gamma(s)) = c + s.$$

Since the Malgrange condition is not affected by a change of origin of \mathbb{R}^n , we can assume that for every small enough positive real number ε there exists a constant A_ε so that

$$(4.3) \quad |\mathbf{x}| \cdot |\nabla \mathbf{t}_M(\mathbf{p})| \geq A_\varepsilon \text{ for all } \mathbf{p} = (\mathbf{x}, t) \in \mathbf{t}_M^{-1}[c - \varepsilon, c + \varepsilon].$$

For $|s| \leq \varepsilon$ we deduce

$$|\gamma(s)| \leq |\gamma(0)| + \int_0^{|s|} \frac{1}{|\nabla \mathbf{t}_M(\gamma(z))|} dz.$$

Combining this latter inequality with Gronwall Lemma provides

$$|\gamma(s)| \leq |\gamma(0)| \cdot \exp\left(\frac{|s|}{A_\varepsilon}\right).$$

Since χ is C^{1+m} in $M \setminus \text{crit}(\mathbf{t}_M)$, the function \mathbf{t}_M is C^{1+m} -trivial at c by the flow of χ with initial conditions along M_c . \square

Definition 4.5. *The set of generalized critical values is defined as*

$$K(\mathbf{t}_M) := K_0(\mathbf{t}_M) \cup K_\infty(\mathbf{t}_M).$$

The Malgrange condition at a regular value c encodes the geometry at infinity of the pencil of nearby fibres. Indeed we have the following

Lemma 4.6. *Let c be a value taken by \mathbf{t}_M which is not a generalized critical value. Let $(\mathbf{p}_k)_k$ be a sequence of points of M converging in \overline{M} to (\mathbf{v}, c) in $\mathbf{S}_\infty^{n-1} \times \{c\} \cap \overline{M}$ while $\mathbf{t}_M(\mathbf{p}_k)$ goes to c . Assume that $\nu_{\mathbf{t}_M}(\mathbf{p}_k)$ converges in \overline{M} to ν in \mathbf{S}^n . Then ν is orthogonal to \mathbf{v} .*

Proof. These limits can be achieved along a C^1 globally definable path $]0, 1[\mapsto M$, $s \mapsto \mathbf{p}(s) = (\mathbf{x}(s), t(s))$ as s goes to 0 with $\lim_0 \mathbf{p} = (\mathbf{v}, c) \in M^\infty$. We choose the parameterization of \mathbf{p} so that $|\mathbf{x}(s)| = s^{-1}$. Let $\mathbf{t}(s) := \mathbf{t}_M(\mathbf{p}(s))$ and so on. Let us write

$$\begin{aligned} \mathbf{t}(s) &= c + As^a + o(s^a) \\ \mathbf{x}(s) &= s^{-1}\mathbf{v} + o(s^{-1}) \\ \nu_{\mathbf{t}_M}(s) &= \nu + s^d\nu_1 + o(s^d) \\ \nu_M^{\mathbf{x}}(s) &= s^e\nu^{\mathbf{x}} + o(s^e) \\ \nu_M^t(s) &= s^f\nu^t + o(s^f) \end{aligned}$$

where $a, d, e, f \in (\mathbb{F}_M)_{\geq 0} \cup \{+\infty\}$ with a, d positive exponents, $\min(e, f) = 0$, and $A \in \mathbb{R}$, $\nu_1 \in \mathbb{R}^n \times \mathbb{R}$, $\nu^{\mathbf{x}} \in \mathbb{R}^n \times 0$, $\nu^t \in \mathbb{R}$ are non-zero vectors whence the corresponding exponent is not ∞ and

$$\nu = -\nu^t \frac{\nu^{\mathbf{x}}}{|\nu^{\mathbf{x}}|} + |\nu^{\mathbf{x}}| \partial_t.$$

We deduce that there exists a continuous definable function $\varphi : (\mathbb{R}_{\geq 0}, 0) \mapsto \mathbb{R}$ with $\varphi(0) = Aa$, such that

$$\mathbf{t}'(s) = s^{a-1}\varphi(s) \quad \text{and} \quad \mathbf{x}'(s) = -s^{-2}\mathbf{v} + o(s^{-2}).$$

Using the Malgrange condition provides

$$1 \geq |\nu_M^{\mathbf{x}}(s)| = |\nabla \mathbf{t}_M(s)| \geq A_c s$$

for some positive constant A_c . Thus $e \leq 1$. Since

$$0 = \langle \nu_M^{\mathbf{x}} + \nu_M^t \partial_t, \mathbf{x}' + \mathbf{t}' \partial_t \rangle = \langle \nu_M^{\mathbf{x}}, \mathbf{x}' \rangle + \nu_M^t \mathbf{t}'$$

we deduce that

$$\nu_M^t \mathbf{t}' = -\langle \nu_M^{\mathbf{x}}, \mathbf{x}' \rangle = s^{e-2}[-\langle \nu^{\mathbf{x}}, \mathbf{v} \rangle + o(1)].$$

From this last equation we deduce that there exists an exponent $e' \geq e$ such that

$$s^{a-1+f} \sim |\nu_M^t \mathbf{t}'| = |\langle \nu_M^{\mathbf{x}}, \mathbf{x}' \rangle| \sim s^{e'-2},$$

so that $\langle \nu^{\mathbf{x}}, \mathbf{v} \rangle = 0$. \square

To conclude this section we introduce a final regularity condition.

Definition 4.7. Let c be a regular value of \mathbf{t}_M taken by \mathbf{t}_M .

The function \mathbf{t}_M is horizontally spherical at c at infinity if for any sequence $(\mathbf{p}_k)_k$ of M converging to $(\mathbf{u}, c) \in M^\infty$, then

$$(4.4) \quad \left\langle \lim_{\infty} \frac{\nu_M^{\mathbf{x}}}{|\nu_M^{\mathbf{x}}|}, \mathbf{u} \right\rangle = 0,$$

where $\lim_{\infty} \frac{\nu_M^{\mathbf{x}}}{|\nu_M^{\mathbf{x}}|}$ means the closed set of all the possible accumulation values, as k goes to infinity, of the unitary vector field of $\frac{\nu_M^{\mathbf{x}}}{|\nu_M^{\mathbf{x}}|}$ along the sequence $(\mathbf{p}_k)_k$.

Note that the following holds true:

Lemma 4.8. The condition of Equation 4.4 is equivalent to

$$\langle \lim_{\infty} \nu_{\mathbf{t}_M}, \mathbf{u} \rangle = 0$$

along any sequence $(\mathbf{p}_k)_k$ of M converging to a point (\mathbf{u}, c) in M^∞ .

Indeed, similarly to what has been done for globally definable functions, we have the following:

Proposition 4.9. Let c be a regular value taken by \mathbf{t}_M . The function \mathbf{t}_M is horizontally spherical at c at infinity if and only if there exists an exponent e_c in $\mathbb{F}_{\mathcal{M}} \cap]-\infty, 1[$ and a positive constant E_c , such that there exist positive real numbers ε and R such that

$$(4.5) \quad (\mathbf{x}, t) \in \mathbf{t}_M^{-1}([c - \varepsilon, c + \varepsilon]) \setminus \mathbf{B}_R \implies |\mathbf{x}| \cdot |\nabla \mathbf{t}_M| \geq E_c |\mathbf{t}_M(\mathbf{x}, t) - c|^{e_c}.$$

Proof. In this globally definable and polynomially bounded context, we can show (as in [d'AcGr2]) that a Bochnak-Łojasiewicz inequality type at the value c not in $K_0(\mathbf{t}_M)$ at infinity holds: there exists a positive constant L_c such that there exist positive real numbers ε and R such that

$$(4.6) \quad (\mathbf{x}, t) \in \mathbf{t}_M^{-1}([c - \varepsilon, c + \varepsilon]) \setminus \mathbf{B}_R \implies |\mathbf{x}| \cdot |\nabla \mathbf{t}_M| \geq L_c |\mathbf{t}_M(\mathbf{x}, t) - c|.$$

1) Assume \mathbf{t}_M is horizontally spherical at c at infinity.

Let $\mathbf{p} :]0, 1[\rightarrow M$ be any continuous globally definable path such that it goes to (\mathbf{u}, c) as s goes to 0. Writing $\mathbf{p} = (\mathbf{x}, t)$ and parameterizing as $|\mathbf{x}(s)| = s^{-1}$, we have

$$\mathbf{p}(s) = (s^{-1}\mathbf{u} + o(s^{-1}), c + As^a + o(s^a))$$

for $A \neq 0$ and $a \in (\mathbb{F}_{\mathcal{M}})_{>0} \cup \{+\infty\}$. The numbers A and a depend on the choice of the path $s \mapsto \mathbf{p}(s)$. We obtain that along \mathbf{p} there exists $a' \leq a$ such that

$$|\mathbf{x}| \cdot |\nabla \mathbf{t}_M| \sim s^{a'}.$$

Note that

$$a' < a \iff \lim_0 \left\langle \nu_{\mathbf{t}_M}, \frac{\mathbf{p}}{|\mathbf{p}|} \right\rangle = 0.$$

In particular the latter equivalence shows that

$$\mathbf{t}_M(\mathbf{x}, t) \rightarrow c \text{ as } \mathbf{x} \rightarrow +\infty \implies \frac{|\mathbf{t}_M(\mathbf{x}, t) - c|}{|\mathbf{x}| \cdot |\nabla \mathbf{t}_M(\mathbf{x}, t)|} \rightarrow 0 \text{ as } \mathbf{x} \rightarrow +\infty.$$

Let ε_0 be a small enough positive number such that $[c - \varepsilon_0, c + \varepsilon_0]$ contains only a single asymptotic critical value: c . Let R_0 be a positive large enough number. Let V_{ε_0, R_0} be the globally definable subset defined as

$$V_{\varepsilon_0, R_0} := \{(\mathbf{x}, t) \in M : |\mathbf{t} - c| \leq \varepsilon_0, |\mathbf{x}| \geq R_0\}.$$

For $R \geq R_0$ let $\mu_0 : [R_0, +\infty[\rightarrow \mathbb{R}$ be defined as

$$\mu_0(R) := \min\{|\mathbf{x}| \cdot |\nabla \mathbf{t}_M(\mathbf{x}, t)| \text{ for } (\mathbf{x}, t) \in V_{\varepsilon_0, R_0} \text{ and } |\mathbf{x}| = R\}.$$

The function μ_0 is globally definable and tends to 0 as R goes to infinity since c is an ACV. If R_0 is large enough, we can write

$$\mu_0(R) = A_0 R^{-a_0} (1 + o(1)) \text{ with } A_0 > 0, a_0 \in (\mathbb{F}_{\mathcal{M}})_{>0}.$$

Let V_0 be the closure of V_{ε_0, R_0} in \overline{M} , thus V_0 is compact in $\overline{\mathbb{R}^n} \times \mathbb{R}$. Let W_0 be the part at infinity of V_0 , that is

$$W_0 := V_0 \cap (\mathbf{S}_{\infty}^{n-1} \times \mathbb{R}).$$

The function

$$\psi_0 : V_0 \setminus W_0 \ni (\mathbf{x}, t) \rightarrow \frac{|\mathbf{t}_M(\mathbf{x}, t) - c|}{|\mathbf{x}| \cdot |\nabla \mathbf{t}_M(\mathbf{x}, t)|}$$

extends continuously and definably over V_0 taking the value 0 along W_0 , by hypothesis of horizontal sphericalness. In the same way, the function

$$\rho_0 : V_0 \setminus W_0 \ni (\mathbf{x}, t) \rightarrow |\mathbf{x}|^{-1}$$

also extends continuously and definably over V_0 taking the value 0 along W_0 . Furthermore we see that

$$\rho_0 = 0 \implies \psi_0 = 0.$$

Thus by a Łojasiewicz argument, there exist a positive exponent b and a positive constant B such that in V_0 the following inequality holds true:

$$\psi_0 \leq B \rho_0^b \iff \psi_0 \leq B |\mathbf{x}|^{-b}.$$

Let μ_1 be the function defined as follows:

$$\mu_1 : V_0 \setminus W_0 \ni (\mathbf{x}, t) \rightarrow \mu_0(|\mathbf{x}|).$$

The function μ_1 is globally definable, continuous and extends continuously to V_0 taking the value 0 along W_0 . Therefore we deduce that in $V_0 \setminus W_0$ we have

$$|\mathbf{t}_M(\mathbf{x}, t) - c| \leq C_0 \cdot \mu_1^{\frac{b}{a_0}} \cdot |\mathbf{x}| \cdot |\nabla \mathbf{t}_M(\mathbf{x}, t)| \leq C_0 \cdot (|\mathbf{x}| \cdot |\nabla \mathbf{t}_M(\mathbf{x}, t)|)^{\frac{b+a_0}{a_0}}$$

where C_0 is a positive constant. This latter inequality provides the announced result.

2) Assume the inequality holds.

Let $\mathbf{p} :]0, 1[\rightarrow M$ be a globally definable continuous path such that $\lim_0 \mathbf{p} = (\mathbf{u}, c)$. Writing $\mathbf{p} = (\mathbf{x}, \mathbf{t})$ and parameterizing as $|\mathbf{x}(s)| = s^{-1}$, we have that

$$\begin{aligned} \mathbf{t}(s) &= c + As^a + o(s^a) \\ \mathbf{p}(s) &= (s^{-1}\mathbf{u} + o(s^{-1}), \mathbf{t}(s)) \in \mathbb{R}^n \times \mathbb{R} \\ \nu_M^{\mathbf{x}}(s) &= s^b \nu + o(s^b) \in \mathbb{R}^n \times 0 \\ \nu_M^{\mathbf{t}}(s) &= s^d (\lambda \nu) + o(s^d) \in \mathbb{R}^n \times 0 \end{aligned}$$

with $A \neq 0$ and $a \in (\mathbb{F}_{\mathcal{M}})_{>0} \cup \{\infty\}$ while $b, d \in (\mathbb{F}_{\mathcal{M}})_{\geq 0}$, $\min(b, d) = 0$, with $\lambda \neq 0$ and $\nu \in \mathbb{R}^n \setminus 0$. Since the path \mathbf{p} lies on M , we know that

$$\langle \nu_M, \mathbf{p}' \rangle = \langle \nu_M^{\mathbf{x}}, \mathbf{x}' \rangle + \nu_M^{\mathbf{t}} \mathbf{t}' = 0$$

from which we deduce

$$(4.7) \quad b - 2 \leq d + a - 1.$$

We want to show that ν is orthogonal to \mathbf{u} , in other words $b < d + a + 1$.

We have the following estimates

$$\begin{aligned} |\nabla \mathbf{t}_M|(s) &= |\nu_M^{\mathbf{x}}|(s) \sim s^b \\ |\mathbf{x}| \cdot |\nabla \mathbf{t}_M|(s) &\sim s^{b-1}. \end{aligned}$$

Using Inequality (4.5), we get

$$(4.8) \quad b - 1 \leq e_c \cdot a < a \text{ and } b < a + 1.$$

Since d is non negative, this yields the orthogonality of \mathbf{u} and ν . \square

5. COMPARING REGULARITY CONDITIONS AND TRIVIALITY

We are working within the context of Section 4.

We have introduced previously three regularity conditions at infinity for the function \mathbf{t}_M . We are going to compare them here.

The hypersurface $M \subset \mathbb{R}^n \times \mathbb{R}$ is the definable family of the hypersurfaces $\{T_t\}_{t \in \mathbb{R}}$ of \mathbb{R}^n and \overline{M} is its closure in $\overline{\mathbb{R}^n} \times \mathbb{R}$. Let M^∞ be the intersection of \overline{M} with the boundary at infinity $\mathbf{S}_\infty^{n-1} \times \mathbb{R}$. By Lemma 4.2, the globally definable function $\mathbf{r}_M : \mathbb{R}^n \times \mathbb{R} \mapsto \mathbb{R}$, defined as $(\mathbf{x}, t) \mapsto |\mathbf{x}|$, is transverse to $M \cap \mathbb{R}^n \times \mathbb{R} \times]-A, A[$ for some positive given A whenever \mathbf{x} is large enough.

In Section 3 was defined

$$\mathcal{C}(M) := \mathcal{X}_{\mathbf{r}_M}^\infty$$

the *space of characteristic covectors of M at infinity*, which is a closed definable subset of $\mathbf{S}_\infty^{n-1} \times \mathbb{R} \times \mathbf{G}^\vee(1, n+1)$.

From Definition 3.3, we also know that: (i) the family $\{T_t\}_{t \in \mathbb{R}}$ is t -equisingular at $\mathbf{p} \in M^\infty$ if

$$(5.1) \quad \mathcal{C}(M)_{\mathbf{p}} \cap (\mathcal{X}_\tau^\infty)_{\mathbf{p}} = \emptyset,$$

where $\tau : \overline{\mathbb{R}^n} \times \mathbb{R} \mapsto \mathbb{R}$ is the projection on the last factor and, (ii) the family $\{T_t\}_{t \in \mathbb{R}}$ is t -equisingular at infinity at c if it is t -equisingular at $\mathbf{p} \in M^\infty$ for all $\mathbf{p} \in M^\infty \cap \tau^{-1}(c)$.

Let $\mathbf{p} = (\mathbf{u}, c) \in M^\infty$. The family $\{T_t\}_{t \in \mathbb{R}}$ is t -equisingular at \mathbf{p} if for any sequence $\mathbf{p}_k = (\mathbf{u}_k, t_k)$ converging to \mathbf{p} such that the sequence of T'_k , the tangent space to the level of \mathbf{r}_M through \mathbf{p}_k , converges to T' , then the latter is not contained in $\mathbb{R}^n \times 0$. This definition is more geometric than the Malgrange condition, which is of interest since we have the following:

Proposition 5.1 (see [DiRuTi] for functions). *If the family $\{T_t\}_{t \in \mathbb{R}}$ is t -equisingular at infinity at c then the function \mathbf{t}_M satisfies the Malgrange condition at c .*

Proof. Suppose that the Malgrange condition is not satisfied at c . There is a globally definable path $\gamma : [1, +\infty[\mapsto M$, $\gamma = (\gamma_{\mathbf{x}}, \gamma_t)$, with $\lim_{\infty} \gamma = \mathbf{p} = (\mathbf{u}, c) \in M^\infty$ and such that

$$(|\mathbf{x}| \cdot |\nabla \mathbf{t}_M|) \circ \gamma \rightarrow 0 \text{ as } r \rightarrow +\infty.$$

Equivalently

$$\left(|\mathbf{x}| \cdot \left| \frac{\partial_{\mathbf{x}} F}{\nabla F} \right| \right) \circ \gamma \rightarrow 0.$$

The following vector V in $\mathbb{R}^n \times \mathbb{R}$:

$$V = \frac{\partial_{\mathbf{x}} F}{|\nabla F|} \begin{pmatrix} \mathbf{x} \\ 0 \end{pmatrix} + \frac{1}{|\nabla F|} \begin{pmatrix} \partial_{\mathbf{x}} F \\ \partial_t F \end{pmatrix}$$

is a normal vector to the level of r_M through the point (\mathbf{x}, t) . We see that $V \circ \gamma \rightarrow \begin{pmatrix} 0 \\ \pm 1 \end{pmatrix}$, since

$$\left(|\mathbf{x}| \cdot \left| \frac{\partial_{\mathbf{x}} F}{\nabla F} \right| \right) \circ \gamma \rightarrow 0, \quad \left| \frac{\partial_{\mathbf{x}} F}{\nabla F} \right| \circ \gamma \rightarrow 0 \quad \text{and} \quad \frac{\partial_t F}{|\nabla F|} \rightarrow \pm 1.$$

This contradicts the observation made just above. \square

Since we just have seen that t -equisingularity at infinity implies the Malgrange condition, we need to check if there is a relation between these and sphericalness at infinity. To this end an obvious corollary of Proposition 4.9 is the following:

Corollary 5.2. *Let c not be a generalized critical value. Then \mathfrak{t}_M is horizontally spherical at c at infinity. In other words t -equisingularity at infinity at c implies horizontal sphericalness at c at infinity.*

Proof. It is just reformulating the fact that Malgrange at c is equivalent to have $e_c \leq 0$ in Equation (4.5). \square

We can now state the last result of this section about local triviality:

Theorem 5.3. *Let c be a value at which \mathfrak{t}_M is horizontally spherical at infinity. Then \mathfrak{t}_M is C^{1+m} is locally trivial at c .*

Proof. Once we have moved the origin of $\mathbb{R}^n \times 0$ so that its value is not c , we just have to integrate the field $\chi = \frac{1}{|\nabla \mathfrak{t}_M|} \nu_{\mathfrak{t}_M}$ as before. Inequality (4.5) now holds in $\mathfrak{t}_M^{-1}[c - \varepsilon, c + \varepsilon] \setminus \mathbf{B}_{R_0}$ for a large positive R_0 . As in [d'AcGr1, d'AcGr2] combining it with Gronwall Lemma will show that any trajectory of χ parameterized over $[0, \varepsilon]$ with initial point in $M_c \cap \mathbf{B}_R$ stays in \mathbf{B}_{KR} for some constant K depending only on c and ε . \square

As a final remark, there are polynomial examples in [d'AcGr1] with regular values which are ACV, but with exponent $e_c < 1$.

6. CURVATURE AND ABSOLUTE CURVATURE OF FAMILIES OF GLOBALLY DEFINABLE HYPERSURFACES

Some of the material presented here can also be found in [Gra1] (or adapted from it).

Let H be a globally definable and oriented hypersurface of \mathbb{R}^n of class C^{1+m} with $m \geq 1$.

Assume that now H is connected and let $\nu_H : H \mapsto \mathbf{S}^{n-1}$ be an orientation. The unitary normal mapping ν_H is globally definable and C^m .

Assume that the maximal rank of $d_{\mathbf{x}} \nu_H$ when \mathbf{x} ranges H is $n - 1$.

There exist finitely many definable disjoint connected open subsets $(U_i)_{i \in I}$ of \mathbf{S}^{n-1} such that

$$\text{clos}(\nu_H(H)) = \cup_{i \in I} \text{clos}(U_i)$$

and for each $i \in I$, the mapping ν_H induces a globally definable finite covering

$$\nu_H : H_i \mapsto U_i$$

where $H_i := \nu_H^{-1}(U_i)$ and such that

$$\dim \nu_H(H \setminus (\cup_{i \in I} H_i)) \leq n - 2.$$

Denoting κ_H the determinant of $d\nu_H$, that is the Gauss-Kronecker curvature of M at the considered point, the total Gauss curvature $K(H)$ of H is defined (if it exists, and it does as we see below) as

$$K(H) := \int_H \kappa_H(\mathbf{x}) d\mathbf{x}.$$

An application of the formula of change of variables gives

$$\int_H \kappa_H(\mathbf{x}) d\mathbf{x} = \sum_{i \in I} (-1)^{d_i} \text{vol}_{n-1}(U_i)$$

for $(-1)^{d_i}$ the degree of the covering mapping $\nu_H|_{H_i} : H_i \mapsto U_i$ for each i .

We introduce another average of curvature, namely the *total absolute curvature* $|K|(H)$ of H defined as

$$|K|(H) := \int_H |\kappa_H(\mathbf{x})| d\mathbf{x}$$

Another application of the formula of change of variables yields,

$$\int_H |\kappa_H(x)| dx = \sum_{i \in I} e_i \cdot \text{vol}_{n-1}(U_i)$$

where e_i is the number of sheets of the covering $\nu_H|_{H_i} : H_i \mapsto U_i$.

The hypothesis on the rank of $d\nu_H$ guarantees that e_i is positive. Otherwise both curvatures are 0.

Returning to the notations and hypotheses of Section 4, the hypersurface M can also be seen as a globally definable family of hypersurfaces $\mathcal{F}_{\mathbf{t}_M} := (T_c)_{c \in \text{Im}(\mathbf{t}_M)}$ of \mathbb{R}^n . We can define the following mapping:

$$\begin{aligned} \mathbf{N} : M \setminus \text{crit}(\mathbf{t}_M) &\mapsto \mathbf{S}^{n-1} \\ (\mathbf{x}, t) &\mapsto \frac{\nu_M^\mathbf{x}}{|\nu_M^\mathbf{x}|} \end{aligned}$$

The mapping N is called *the Gauss mapping of the family* $\mathcal{F}_{\mathbf{t}_M}$. It is globally definable and C^{1+m} . The restriction of $N|_{T_c}$ is denoted \mathbf{N}_c , so that the family of mappings $(\mathbf{N}_c)_{c \in \text{Im}(\mathbf{t}_M) \setminus K_0(\mathbf{t}_M)}$ is globally definable, where $\text{Im}(\mathbf{t}_M)$ is the image of the function \mathbf{t}_M . Let κ_c be the Gauss-Kronecker curvature of T_c . Thus we can define two functions

$$\begin{aligned} K : \text{Im}(\mathbf{t}_M) \setminus K_0(\mathbf{t}_M) &\rightarrow \mathbb{R} \\ c &\mapsto K(c) := \int_{T_c} \kappa_c(\mathbf{p}) d\mathbf{p}, \\ |K| : \text{Im}(\mathbf{t}_M) \setminus K_0(\mathbf{t}_M) &\rightarrow \mathbb{R} \\ c &\mapsto |K|(c) := \int_{T_c} |\kappa_c|(\mathbf{p}) d\mathbf{p}. \end{aligned}$$

The introductory material of this section guarantees that both functions are well defined. The paper [Gra1] has dealt with the case where M is a graph. We can state now the result of this section:

Theorem 6.1. (i) *There are finitely many values in $\text{Im}(\mathbf{t}_M) \setminus K_0(\mathbf{t}_M)$ at which the function $t \mapsto K(t)$ is not continuous*

(ii) *There are finitely many values in $\text{Im}(\mathbf{t}_M) \setminus K_0(\mathbf{t}_M)$ at which the function $t \mapsto |K|(t)$ is not continuous*

(iii) *If $|K|$ is continuous at c , so is K .*

Sketch of Proof. It is a very similar proof to that of [Gra1, Sections 4,5,6].

Let us consider the following globally definable and C^{1+m} mapping

$$\begin{aligned} \Psi : M &\mapsto \mathbf{S}^{n-1} \times \mathbb{R} \\ \mathbf{p} &\mapsto (\mathbf{N}(\mathbf{p}), \mathbf{t}_M(\mathbf{p})). \end{aligned}$$

It is a local diffeomorphism at any point of $M \setminus \text{crit}(\Psi)$. Let $\Delta := \Psi(\text{crit}(\Psi))$ which is definable, closed and of dimension lower than or equal to $n - 1$. Let $\mathcal{U} := (\mathbf{S}^{n-1} \times \mathbb{R}) \setminus \Delta$.

There exists an integer number p_M such that for any $(\mathbf{u}, t) \in \mathcal{U}$ the fibre $\Psi^{-1}(\mathbf{u}, t)$ has at most p_M points. For any point (\mathbf{u}, t) in \mathcal{U} the degree $\delta(\mathbf{u}, t)$ of Ψ at (\mathbf{u}, t) may range from $-p_M$ to p_M . In particular the function $(\mathbf{u}, t) \mapsto \delta(\mathbf{u}, t)$ is definable and

$$\delta(\mathbf{u}, t) = \deg_{\mathbf{u}} \mathbf{N}_t.$$

We define the following subsets

$$\begin{aligned} \mathcal{U}_k &:= \{(\mathbf{u}, t) \in \mathcal{U} : \#\Psi^{-1}(\mathbf{u}, t) = k\} \\ U_t &:= \{\mathbf{u} \in \mathbf{S}^{n-1} : (\mathbf{u}, t) \in \mathcal{U}\} = \mathbf{N}_t(T_t \setminus \text{crit}(\mathbf{N}_t)) \\ U_{t,k} &:= \{\mathbf{u} \in \mathbf{S}^{n-1} : (\mathbf{u}, t) \in \mathcal{U}_k\}. \end{aligned}$$

The subsets U_t and $U_{t,k}$ are open, and we obtain finitely many globally definable families $(U_t)_{t \in \text{Im}(\mathbf{t}_M) \setminus K_0(\mathbf{t}_M)}$ and $(U_{t,k})_{t \in \text{Im}(\mathbf{t}_M) \setminus K_0(\mathbf{t}_M)}$.

Note that $U_t = \cup_k U_{t,k}$ and since the function $\mathbf{u} \mapsto \deg_{\mathbf{u}} \mathbf{N}_t$ is definable, it is constant on each connected component of U_t .

Let c be a regular value of \mathbf{t}_M . Since Hausdorff limits of closed definable subsets of a given compact space exist, we can set

$$\begin{aligned} \mathcal{V}_c^+ &:= \lim_{t \rightarrow c, t > c} \text{clos}(U_t) & \text{and} & & \mathcal{V}_{c,k}^+ &:= \lim_{t \rightarrow c, t > c} \text{clos}(U_{t,k}) \\ \mathcal{V}_c^- &:= \lim_{t \rightarrow c, t < c} \text{clos}(U_t) & \text{and} & & \mathcal{V}_{c,k}^- &:= \lim_{t \rightarrow c, t < c} \text{clos}(U_{t,k}). \end{aligned}$$

Let V_1, \dots, V_{d_c} be the connected components of U_c . For each $i = 1, \dots, s$, let k_i be the integer number such that $V_i \subset U_{c,k_i}$. For each $i = 1, \dots, d_c$ there exists $l_i^+ \geq k_i$ and $l_i^- \geq k_i$ such that

$$V_i \subset \mathcal{V}_{c,l_i^-}^- \quad \text{and} \quad V_i \subset \mathcal{V}_{c,l_i^+}^+.$$

In particular we deduce that for each i

$$\text{vol}_{n-1}(V_i) \leq \min\{\text{vol}_{n-1}(\mathcal{V}_{c,l_i^-}^-), \text{vol}_{n-1}(\mathcal{V}_{c,l_i^+}^+)\}.$$

Let δ_i be the degree of \mathbf{N}_c at any point of V_i . We find

$$K(c) = \sum_{i=1}^{d_c} \delta_i \cdot \text{vol}_{n-1}(V_i) \quad \text{and} \quad |K|(s) = \sum_{i=1}^{d_c} k_i \cdot \text{vol}_{n-1}(V_i).$$

From the previous arguments we get that each following limit exists

$$K_c^+ := \lim_{t \rightarrow c, t > c} K(t), \quad K_c^- := \lim_{t \rightarrow c, t < c} K(t), \quad |K|_c^+ := \lim_{t \rightarrow c, t > c} |K|(t), \quad |K|_c^- := \lim_{t \rightarrow c, t < c} |K|(t),$$

and we obviously get

$$|K|(c) \leq \min(|K|_c^-, |K|_c^+).$$

The rest of the proof follows from the following arguments: Assume that each U_t has d_t connected $V_{t,1}, \dots, V_{t,d_t}$. Each such connected component $V_{t,i}$ lies in $U_{t,k_i(t)}$ with $k_i(t) \leq k_j(t)$ if and only if $i \leq j$. Moreover the degree of \mathbf{N}_t at any point of $V_{t,i}$ is constant and equal to $\delta_i(t)$. These comes from properties of Ψ and \mathcal{U} . From here we deduce that there exists a finite subset Z of \mathbb{R} such that for any J connected component of $(\mathbb{R} \setminus Z) \cap \text{Im}(\mathbf{t}_M)$, the numbers $d_t, k_i(t), \delta_i(t)$ are independent of t in J . Moreover each function $t \mapsto \text{vol}_{n-1}(V_{t,i})$ is continuous over J . \square

7. MORE ON REGULARITY AT INFINITY

Let $\mathbf{N} : M \setminus \text{crit}(\mathbf{t}_M) \mapsto \mathbf{S}^{n-1}$ be the Gauss mapping of the family of the regular levels of \mathbf{t}_M . Similarly to the conormal geometry at infinity (in $\mathbb{R}^n \times \mathbb{R}$) of the function \mathbf{t}_M , we are interested in the limits of \mathbf{N} at infinity (in \mathbb{R}^n).

Let $\Gamma(\mathbf{N})$, contained in $M \times \mathbf{S}^{n-1}$, be the graph of \mathbf{N} , let $\overline{\Gamma(\mathbf{N})}$ be its closure in $\overline{\mathbb{R}^n} \times \mathbb{R} \times \mathbf{S}^{n-1}$ and $\overline{\mathbf{N}} : \overline{\Gamma(\mathbf{N})} \mapsto \mathbf{S}^{n-1}$ be the projection onto \mathbf{S}^{n-1} , so that we can think of it as the extension by continuity of \mathbf{N} to $\overline{\Gamma(\mathbf{N})}$.

The closed definable subset $T_{c,+}^\infty$ is defined as

$$T_{c,+}^\infty := \{\mathbf{u} \in \mathbf{S}^{n-1} : \exists (\mathbf{p}_k)_k \in M \text{ such that } \lim_{\infty} \mathbf{p}_k = (\mathbf{u}, c)\}.$$

Let $V_c^\infty := \overline{\mathbf{N}}(\pi^{-1}(T_{c,+}^\infty \times \{c\}))$, in other words it is the definable closed subset

$$V_c^\infty = \{\mathbf{v} \in \mathbf{S}^{n-1} : \exists ((\mathbf{x}_k, \tau_k))_k \in M \text{ such that } \mathbf{x}_k \rightarrow \infty, \tau_k \rightarrow c, \mathbf{N}(\mathbf{x}_k, \tau_k) \rightarrow \mathbf{v}\},$$

corresponding to all the limits at infinity of normals to the hypersurfaces $(T_t)_t$ as t tends to c .

For each $\mathbf{u} \in \mathbf{S}_{\infty}^{n-1}$, let $V_{c,\mathbf{u}}^{\infty} := V_c^{\infty} \cap \overline{\mathbf{N}}(\pi^{-1}(\mathbf{u}) \times \{c\})$, that is

$$V_{c,\mathbf{u}}^{\infty} = \left\{ \mathbf{v} \in \mathbf{S}^{n-1} : \exists ((\mathbf{x}_k, \tau_k))_k \in M \text{ such that } \mathbf{x}_k \rightarrow \infty, \tau_k \rightarrow c, \frac{\mathbf{x}_k}{|\mathbf{x}_k|} \rightarrow \mathbf{u}, \mathbf{N}(\mathbf{x}_k, \tau_k) \rightarrow \mathbf{v} \right\}.$$

Note that whenever \mathbf{u} does not belong to $T_{c,+}^{\infty}$ we find that $V_{c,\mathbf{u}}^{\infty}$ is empty.

A very rigid consequence of \mathbf{t}_M being horizontally spherical at c at infinity is the following:

Lemma 7.1. *Let c be a regular value taken by \mathbf{t}_M at which it is horizontally spherical at infinity. Then each \mathbf{u} in $T_{c,+}^{\infty}$ and each \mathbf{v} in $V_{c,\mathbf{u}}^{\infty}$ are orthogonal.*

Proof. Obvious from the definition of the horizontal sphericalness. □

Let c be a regular value taken by \mathbf{t}_M at which it is horizontally spherical at infinity. Let ε be a positive real number such that for each $t \in [c - \varepsilon, c + \varepsilon]$ the function \mathbf{t}_M is horizontally spherical at t at infinity. Let $T_{c,\varepsilon} := \mathbf{t}_M^{-1}([c + \varepsilon, c - \varepsilon])$.

We find that for each for η in $]e_c, 1[$, there exists a positive real number R such that for every (\mathbf{x}, t) belonging to $T_{c,\varepsilon} \setminus \text{clos}(\mathbf{B}_R)$, we have

$$(7.1) \quad |\mathbf{x}| \cdot |\nabla \mathbf{t}_M(\mathbf{x}, t)| \geq |t - c|^{\eta}.$$

Let ξ be the following definable vector field

$$\xi := \frac{1}{|\nabla \mathbf{t}_M|} \frac{\nu_{\mathbf{t}_M}^{\mathbf{S}}}{|\nu_{\mathbf{t}_M}^{\mathbf{S}}|}, \text{ for } |(\mathbf{x}, t)| \geq R \gg 1, \text{ and } |t - c| \leq \varepsilon.$$

It is definable and C^{1+m} , non vanishing, tangent to the Euclidean spheres. The flow of the differential equation

$$\dot{\mathbf{p}}(t) = \xi(\mathbf{p}(t)) \text{ and } \xi(0) \in T_c \times \{c\} \setminus \mathbf{B}_R$$

induces a C^{1+m} diffeomorphism $(T_c \times \{c\} \setminus \mathbf{B}_R) \times [-\varepsilon, \varepsilon] \mapsto T_{c,\varepsilon} \setminus \mathbf{B}_R$.

Using Inequality (7.1) we deduce that the length $l(\mathbf{p}_0, \mathbf{p}_t)$ of the trajectory of ξ between the point \mathbf{p}_0 of $T_c \times \{c\} \setminus \mathbf{B}_R$ and \mathbf{p}_t , point reached after time t , is bounded as

$$(7.2) \quad l(\mathbf{p}_0, \mathbf{p}_t) \leq |\mathbf{p}_0| \left(\frac{t^{1-\eta}}{1-\eta} \right).$$

Inequality (7.2) implies that the angle $\theta(t)$ between the vector \mathbf{p}_t and \mathbf{p}_0 tends to 0 as t goes to 0. This proves the following:

Lemma 7.2. *Let c be a regular value taken by the function \mathbf{t}_M at which it is horizontally spherical at infinity. Then $T_c^{\infty} = T_{c,+}^{\infty}$, thus $T_{c,+}^{\infty}$ is of dimension at most $n - 2$.*

8. MAIN RESULT

Our main result Theorem 8.1 presented in this section is a consequence of results of equisingularity theory and of our context.

Theorem 8.1. *Let $F : \mathbb{R}^n \times \mathbb{R} \mapsto \mathbb{R}$ be a C^{2+m} globally definable function over a polynomially bounded o -minimal structure, for a non negative integer number m . Assuming that 0 is regular value of F , let M be the level $\{F = 0\}$. Let \mathbf{t}_M be the projection of M onto \mathbb{R} .*

Let c be a regular value taken by \mathbf{t}_M at which it is horizontally spherical at infinity. Then the total absolute curvature function $t \mapsto |K|(t)$ is continuous at c . Consequently the total curvature function $t \mapsto K(t)$ is continuous at c .

It is a straightforward consequence of the following

Lemma 8.2. *Under the hypotheses of Theorem 8.1, we find*

$$\dim V_c^{\infty} \leq n - 2.$$

Let us show the main result.

Proof of the main result. Let $|K| : t \mapsto |K|(t)$ be the total absolute curvature function of the family of hypersurfaces $(T_t)_t$. By Lemma 8.2 we find that V_c^∞ has $(n-1)$ -dimensional volume zero. Following [Gra1, Proposition 6.8], we deduce there is no accumulation of curvature at infinity at c . In other words the function $|K|$ is continuous at c , and so is K by point (iii) of Theorem 6.1. \square

Before going into the proof of Lemma 8.2, we observe that it states that there is no accumulation of curvature at infinity nearby the level c , or equivalently there are no half-branch at infinity of the generic polar curve along which the function \mathbf{t}_M tends to c (see [Tib1, Gra2] for local triviality results with a similar flavor).

Proof of Lemma 8.2. Let \mathbf{v} be a limit of normal direction lying in V_c^∞ . By the Curve Selection Lemma we can find a globally definable continuous path, going to infinity, along which this limit is reached: there exists such a path γ such that

$$\mathbf{N} \circ \gamma \rightarrow \mathbf{v} \quad \text{and} \quad \mathbf{t}_M \circ \gamma \rightarrow c.$$

In particular there exists a positive exponent α , in the field of exponents \mathbb{F}_M of the structure \mathcal{M} such that

$$\frac{\mathbf{t}_M \circ \gamma - c}{|\gamma|^\alpha} \rightarrow a \in \mathbb{R}^*.$$

In other words there exists a positive exponent e such that the germ at infinity of γ lies in

$$\mathcal{H}_e := \{\mathbf{p} \in T_{c,\varepsilon} \setminus \mathbf{B}_R : |\mathbf{t}_M(\mathbf{p}) - c| \leq |\mathbf{p}|^{-e}\}.$$

If the exponent e belongs to F_M , then \mathcal{H}_e is globally definable and so is its closure $\overline{\mathcal{H}_e}$ in $\overline{\mathbb{R}^n} \times \mathbb{R}$. Let us define

$$V_c^{\infty,e} := \{\mathbf{v} \in \mathbf{S}^{n-1} : \exists (\mathbf{p}_k)_k \in \mathcal{H}_e \text{ such that } \mathbf{N}(\mathbf{p}_k) \rightarrow \mathbf{v}\}$$

which is a closed definable subset of \mathbf{S}^{n-1} contained in V_c^∞ whenever e lies in \mathbb{F}_M .

Let \mathcal{H}_e^∞ be the intersection $\overline{\mathcal{H}_e} \cap \mathbf{S}_\infty^{n-1} \times \{c\}$. The function \mathbf{t}_M extends continuously and definably to $\overline{\mathcal{H}_e}$ taking the value c along \mathcal{H}_e^∞ . Let \mathbf{t}_e be the restriction of this extension to $\overline{\mathcal{H}_e} \setminus \mathbf{B}_R$.

According to [Bek, Loi], we can stratify the pair $(\mathbf{t}_e, \overline{\mathcal{H}_e} \setminus \mathbf{B}_R)$ with Thom's condition. Furthermore we can require that $X := \mathcal{H}_e \setminus \mathbf{B}_R$ and $Y := \mathcal{H}_e^\infty$ are union of strata.

Suppose first that X and Y are strata. The dimension of Y is $d \leq n-2$ since \mathcal{H}_e^∞ is contained in $T_{c,+}^\infty$, thus of dimension lower than or equal to $n-2$ by Lemma 7.2. Let $\mathbf{p} = (\mathbf{u}, c)$ be a point of Y and let $T := T_{\mathbf{p}}Y$ which is contained in $\mathbb{R}^n \times 0$. Note that T and \mathbf{u} are orthogonal.

Let \mathbf{v} be a limit of the normal \mathbf{N} at infinity at \mathbf{u} taken into \mathcal{H}_e along a path γ . We will show that \mathbf{v} and $T \oplus \mathbb{R}\mathbf{u}$ are orthogonal. We recall that $\nu_M = \nu_M^x + \nu_M^t \partial_t$. Let ν be the limit of ν_M along γ as γ goes to infinity and let η be the limit of $\nu_{\mathbf{t}_M}$. Writing ν as (ν^x, ν^t) in $\mathbb{R}^n \times \mathbb{R}$, we have

$$\mathbf{v} = \frac{\nu^x}{|\nu^x|} \quad \text{and} \quad \eta = -\nu^t \mathbf{v} + |\nu^x|^2 \partial_t.$$

Thom's condition implies that η and T are orthogonal. Moreover, by horizontal spherical-ness at infinity, η and \mathbf{u} are also orthogonal, therefore η and $T \oplus \mathbb{R}\mathbf{u}$ are orthogonal too. Hence, if $\nu^t \neq 0$, then \mathbf{v} is orthogonal to $T \oplus \mathbb{R}\mathbf{u}$ since $T \oplus \mathbb{R}\mathbf{u}$ is contained in $\mathbb{R}^n \times 0$. If $\nu^t = 0$ then $\mathbf{v} = \nu$. Using the arguments of the proof of Lemma 4.2, we see that \mathbf{u} and ν are orthogonal. By Whitney's condition (a), we know that $T_{\mathbf{p}}Y$ is a subspace of $\lim_{\infty} T_\gamma M$ and so $T_{\mathbf{p}}Y$ and ν are orthogonal. Hence we conclude that $\mathbf{v} = \nu$ is orthogonal to $T_{\mathbf{p}}Y \oplus \mathbb{R}\mathbf{u}$.

Let $V_{c,\mathbf{u}}^{\infty,e} := V_c^{\infty,e} \cap V_{c,\mathbf{u}}^\infty$. We have proved that $\dim V_{c,\mathbf{u}}^{\infty,e} \leq (n-1) - (d+1) = n-d-2$, and thus $\dim V_c^{\infty,e} \leq n-2$.

In the general case the only thing to check is that whenever X contains a (globally definable) stratum S of dimension s at most $n-1$, then its contribution to V_c^∞ is at most of dimension $n-2$. But this is so since the graph of $\mathbf{N}|_S$ is of dimension s , so that its limits at infinity

$$\{\mathbf{v} \in \mathbf{S}^{n-1} : \exists S \ni (\mathbf{x}_k, \tau_k)_k \text{ such that } \tau_k \rightarrow c, \mathbf{N}(\mathbf{x}_k, \tau_k) \rightarrow \mathbf{v}\} \subset V_c^\infty$$

have dimension at most $s - 1 \leq n - 2$.

We conclude that $V_c^{\infty, e}$ has dimension lower than or equal to $n - 2$ for any exponent e of $\mathbb{F}_{\mathcal{M}}$.

Since any limit \mathbf{v} of V_c^∞ belongs to some $V_c^{\infty, e}$ for some e in $\mathbb{F}_{\mathcal{M}}$, and since the family $(V_c^{\infty, e})_{e \in (\mathbb{F}_{\mathcal{M}})_{>0}}$ is increasing as e goes to 0, we get that V_c^∞ is the Hausdorff limit at $e = 0$ of $V_c^{\infty, e}$, thus has dimension lower than or equal to $n - 2$. \square

We conclude with an interesting observation. For this purpose we need a few more preparations. Let c be regular value taken by \mathbf{t}_M . Let ε be a positive number such that $[c - \varepsilon, c + \varepsilon]$ consists only of regular values. Let Z be a connected component of $\mathbf{t}_M^{-1}(]c - \varepsilon, c + \varepsilon[)$. Let us consider now \mathbf{t}_Z the restriction of \mathbf{t}_M to Z . Let $Z_t \times \{t\} := \mathbf{t}_Z^{-1}(t) = M_t \cap Z$. Let $K_Z(t) := \int_{Z_t} \kappa$ and $|K|_Z(t) := \int_{Z_t} |\kappa|$ for t in $]c - \varepsilon, c + \varepsilon[$. Then we actually have showed the following:

Theorem 8.3. *Under the above hypotheses, assume furthermore that \mathbf{t}_Z is horizontally spherical at infinity at c . Then the functions K_Z and $|K|_Z$ are continuous at c .*

To rephrase informally Theorem 8.3, the continuity of $t \mapsto |K|(t)$ nearby the value c at which the function \mathbf{t}_M is horizontally spherical at infinity, is equivalent to the continuity nearby c of each function $t \mapsto |K|_Z(t)$ for each connected component Z of $\mathbf{t}_M^{-1}(]c - \varepsilon, c + \varepsilon[)$.

9. THE SPECIAL CASE OF FUNCTIONS

We treat here briefly the case of functions which is a special case of the context presented here. The continuity of curvatures is the same property but the regularity conditions are a little bit different.

Let $f : \mathbb{R}^n \mapsto \mathbb{R}$ be a C^{2+m} , with non-negative m , globally definable function. We denote the level $f^{-1}(t)$ by F_t and its closure in the spherical compactification by \overline{F}_t . Its intersection with the sphere at infinity \mathbf{S}_∞^{n-1} will be denoted F_t^∞ . Let ν_f be the unitary gradient field $\frac{\nabla f}{|\nabla f|}$.

The function f satisfies *Malgrange condition* at c if there are positive constants R, ε, A such that

$$|\mathbf{x}| > R, |f(\mathbf{x}) - c| < \varepsilon \implies |\mathbf{x}| \cdot |\nabla f(\mathbf{x})| \geq A|f(\mathbf{x}) - c|.$$

We would like to introduce what the analogue of horizontal spherical-ness in this context would be. The function f is *spherical at the regular value c at infinity* if along any sequence of points $(\mathbf{x}_k)_k$ of \mathbb{R}^n such that $|\mathbf{x}_k|$ goes to ∞ and $f(\mathbf{x}_k)$ goes to c , we have

$$\left\langle \lim_{\infty} \nu_f(\mathbf{x}_k), \lim_{\infty} \frac{\mathbf{x}_k}{|\mathbf{x}_k|} \right\rangle = 0,$$

whenever each limit exists.

This condition is equivalent to the following result already proved in [d'AcGr1, d'AcGr2] which justified the introduction for families of the notion of horizontal spherical-ness at infinity.

Theorem 9.1 ([d'AcGr1, d'AcGr2]). *Let c be a regular value of f taken by f . The function f is spherical at infinity at c if and only if there exists an exponent e_c in $\mathbb{F}_{\mathcal{M}} \cap]-\infty, 1[$ and a positive constant E_c such that*

$$|\mathbf{x}| \gg 1, |f(\mathbf{x}) - c| \ll 1 \implies |\mathbf{x}| \cdot |\nabla f(\mathbf{x})| \geq E_c |f(\mathbf{x}) - c|^{e_c}.$$

It is well known that t -regularity is equivalent to Malgrange [DiRuTi] (their proof goes through the globally definable context) and that Malgrange is equivalent to requiring having $e_c \leq 0$, thus spherical-ness at infinity.

Let $K(t)$ be the total Gauss-Kronecker curvature of F_t and $|K|(t)$ be the total absolute Gauss-Kronecker curvature of F_t . In the context of functions what we have proved is the following:

Theorem 9.2. *Let $f : \mathbb{R}^n \mapsto \mathbb{R}$ be a globally definable C^{2+m} function for some non-negative integer m . Let c be a regular value at which the function is spherical at infinity.*

(1) *Then the function $t \mapsto |K|(t)$ is continuous at c , and thus so is $t \mapsto K(t)$.*

(2) *As for Theorem 8.3, for any connected component Z of $f^{-1}[c - \varepsilon, c + \varepsilon]$ [for positive ε small enough, the function $t \mapsto |K|_Z(t)$ is continuous at c , and thus so is $t \mapsto K_Z(t)$.*

Let us end with a last result on equisingularity of the family of fibres of a function.

Corollary 9.3. *Let $f : \mathbb{R}^n \mapsto \mathbb{R}$ be a globally definable C^{2+m} function for some non-negative integer m . Let c be a regular value at which the function is spherical at infinity.*

If n is odd then the following function is continuous at c

$$t \mapsto \int_{\mathbf{G}(n-1,n)} \chi(f^{-1}(t) \cap H) dH.$$

If n is even then the following function is continuous at c

$$t \mapsto \int_{\mathbf{G}(n-1,n)} [\chi(\{f \geq t\} \cap H) - \chi(\{f \leq t\} \cap H)] dH.$$

Proof. By Theorem 9.2, we know that the function $t \mapsto K(t)$ is continuous at c . Then we apply Theorem 4.5 in [Dut1]. If n is odd, the result is clear because the function $t \mapsto \chi(f^{-1}(t))$ is constant in a neighborhood of c . If n is even, it is enough to prove that the functions $t \mapsto \chi(\{f \geq t\})$ and $t \mapsto \chi(\{f \leq t\})$ are constant in a neighborhood of c . By the Mayer-Vietoris sequence, if $t > c$ then we have

$$\chi(\{f \geq c\}) = \chi(\{f \geq t\}) + \chi(\{c \leq f \leq t\}) - \chi(f^{-1}(t)).$$

So if t is close enough to c then $\chi(\{f \geq c\}) = \chi(\{f \geq t\})$, for f is a fibration over $[c, t]$. Similarly we can show that $\chi(\{f \leq c\}) = \chi(\{f \leq t\})$ for $t > c$ close enough to c . The same argument works for $t < c$. \square

REFERENCES

- [Bek] K. BEKKA, *Regular stratification of subanalytic sets*, Bull. London Math. Soc. **25** (1993), 7–16.
- [BrSp] J. BRIANÇON & J.-P. SPEDER, *Les conditions de Whitney impliquent “ μ^* -constant”*, Ann. Inst. Fourier **26** (1976), **2**, xi, 153–163.
- [Com] G. COMTE, *Équisingularité réelle : nombres de Lelong et images polaires*, Ann. Sci. École Norm. Supér. (4) **33** (2000), no. 6, 757–788.
- [ComMe] G. COMTE & M. MERLE, *Équisingularité réelle II : invariants locaux et conditions de régularité*, Ann. Sci. Éc. Norm. Supér. (4) **41** (2008), no. 2, 221–269.
- [CosPe] M. COSTE & M.J. DE LA PUENTE, *Atypical values at infinity of a polynomial function on the real plane: an erratum, and an algorithmic criterion*, J. Pure Appl. Algebra **162** (2001), no. 1, 23–35.
- [d’Ac1] D. D’ACUNTO, *Valeurs critiques asymptotiques d’une fonction définissable dans une structure o-minimale*, Ann. Polon. Math. **75** (2000), no. 1, 35–45.
- [d’Ac2] D. D’ACUNTO, *Sur la topologie des fibres d’une fonction définissable dans une structure o-minimale*, C. R. Acad. Sci. Paris, Sér. I **337** (2003) no. 5, 327–330.
- [d’AcGr1] D. D’ACUNTO & V. GRANDJEAN, *On gradient at infinity of semialgebraic functions*, Ann. Pol. Math., **87**, (2005) 39–49.
- [d’AcGr2] D. D’ACUNTO & V. GRANDJEAN, *On gradient at infinity of tame functions*, Rev. Mat. Complut., **18**, no.2, (2005) 493–501.
- [Dra] R.N. DRAPER, *Intersection theory in analytic geometry*, Math. Ann. **180** (1969) 175–204.
- [Dut1] N. DUTERTRE, *Geometrical and topological properties of real polynomial fibres*, Geometriae Dedicata **105**, no. 1 (2004), 43–59.
- [Dut2] N. DUTERTRE, *Lipschitz-Killing curvatures and polar images*, Advances in Geometry, <https://doi.org/10.1515/advgeom-2018-0019>.
- [vdD] L. VAN DEN DRIES, *Tame Topology and o-Minimal Structures*, London Math. Soc. Lecture Note Ser. 248, Cambridge Univ. Press, 1988.
- [vdDM] L. VAN DEN DRIES & C. MILLER, *Geometric categories and o-minimal structures*, Duke Math. J. **84** (1996), 497–540.
- [DiRuTi] L.R.G. DIAS & M.A.S. RUAS & M. TIBAR, *Regularity at infinity of real mappings and a Morse-Sard theorem*, J. Topology **5** (2012), no. 2, 323–340.
- [Gaf1] T. GAFFNEY, *Multiplicities and equisingularity of ICIS germs*, Invent. Math. **123** (1996), no. 2, 209–220.

- [Gaf2] T. GAFFNEY, *Polar multiplicities and equisingularity of map germs*, *Topology* **32** (1993), no. 1, 185–223.
- [Gra1] V. GRANDJEAN, *On the total curvatures of a tame function*, *Bull. Braz. Math. Soc.* **39** (2008), no. 4, 515–535.
- [Gra2] V. GRANDJEAN, *Tame functions with strongly isolated singularities at infinity: a tame version of a Parusinski’s theorem*, *Geom. Dedicata* **140** (2009), 1–17.
- [HL] H. V. HÀ & D. T. LÊ, *Sur la topologie des polynômes complexes*, *Acta Math. Vietnam.* **9** (1984) 21–32.
- [Hir] H. HIRONAKA, *Normal cones in analytic Whitney stratifications*, *Inst. Hautes Études Sci. Publ. Math.* **36** (1969), 127–138.
- [JoTi] C. JOIȚA & M. TIBĂR, *Bifurcation values of families of real curves*, *Proc. Roy. Soc. Edinburgh Sect. A*, **147** (2017), no. 6, 1233–1242.
- [Kur] K. KURDYKA, *On gradients of functions definable in o-minimal structures*, *Ann. Inst. Fourier* **48**, no. 3 (1998), 769–784.
- [La] R. LANGEVIN, *Courbure et singularités complexes*, *Comment. Math. Helv.* **54** (1979) 6–16.
- [Loe] F. LOESER, *Formules intégrales pour certains invariants locaux des espaces analytiques complexes*, *Comment. Math. Helv.* **59** (1984), no. 2, 204–225.
- [Loi] T. L. LOI, *Verdier stratifications and Thom stratifications in o-minimal structures* *Illinois J. Math* **42**, no. 2 (1998), 347–356.
- [LoZa] T. L. LOI & A. ZAHARIA, *Bifurcation sets of functions definable in o-minimal structures*, *Illinois J. Math.* **42**, no. 3, (1998), 449–457.
- [NgVa] X.V.N. NGUYEN & G. VALETTE, *Whitney stratifications and the continuity of local Lipschitz Killing curvatures*, *Annales de l’Institut Fourier*, **68**, no. 5 (2018), 2253–2276.
- [NeZa] A. NEMETHI & A. ZAHARIA, *Milnor fibration at infinity*, *Indag. Mathem. N.S.* 3 (3), (1992) 323–335.
- [Par] A. PARUSIŃSKI, *On the bifurcation set of complex polynomial with isolated singularities at infinity*, *Compositio Mathematica* 97 (1995) 369–384.
- [SiTi1] D. SIERSMA & M. TIBĂR, *Singularities at infinity and their vanishing cycles*, *Duke Math. J.* **80**, No.3 (1995) 771–783.
- [SiTi2] D. SIERSMA & M. TIBĂR, *The Gauss-Bonnet defect of complex affine hypersurfaces*, *Bull. Sci. Math.* **130**, no. 2 (2006) 110–122.
- [Tei1] B. TEISSIER, *Cycles évanescents, sections planes et conditions de Whitney*, *Singularités à Cargèse (Rencontre Singularités Géom. Anal., Inst. Études Sci., Cargèse, 1972)*, pp. 285–362, *Astérisque*, **7 et 8**, Soc. Math. France, Paris, 1973.
- [Tei2] B. TEISSIER, *Variétés polaires, II. Multiplicités polaires, sections planes, et conditions de Whitney*, *Algebraic geometry (La Rábida, 1981)*, 314–491, *Lecture Notes in Math.*, **961**, Springer, Berlin, 1982.
- [Tho] R. THOM, *Ensembles et morphismes stratifiés*, *bul. Amer. Math. Soc.* **75** (1969) 240–282.
- [Tib1] M. TIBĂR, *Asymptotic equisingularity and topology of complex hypersurfaces* *Int. Math. Res. Not.* No.18, (1998) 979–990.
- [Tib2] M. TIBĂR, *Regularity at infinity of real and complex polynomial functions*, *Singularity Theory*, Edited by Bill Bruce & David Mond, *LMS Lecture Notes*, 263, Cambridge University Press, (1999), 249–264.
- [TiZa] M. TIBĂR & A. ZAHARIA, *Asymptotic behaviour of families of real curves*, *Manuscripta Math.* 99, no.3 (1999) 383–393.
- [Val] G. VALETTE, *Volume, Whitney conditions and Lelong number*, *Ann. Polon. Math.* **93** (2008), no. 1, 1–16.

LABORATOIRE ANGEVIN DE RECHERCHE EN MATHÉMATIQUES, LAREMA, UMR6093, CNRS, UNIV. ANGERS, SFR MATH-STIC, 2 Bd LAVOISIER 49045 ANGERS CEDEX 01, FRANCE.

E-mail address: nicolas.dutertre@univ-angers.fr

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO CEARÁ (UFC), CAMPUS DO PICI, BLOCO 914, CEP. 60455-760. FORTALEZA-CE, BRASIL

E-mail address: vgrandjean@mat.ufc.br