



Is there a link between acetylcholinesterase, behaviour and density populations of the ragworm *Hediste diversicolor*?

Isabelle Métais, Amélie Châtel, Mohammed Mouloud, Hanane Perrein-Ettajani, Mélanie Bruneau, Patrick Gillet, Nisrine Jrad, Catherine Mouneyrac

► To cite this version:

Isabelle Métais, Amélie Châtel, Mohammed Mouloud, Hanane Perrein-Ettajani, Mélanie Bruneau, et al.. Is there a link between acetylcholinesterase, behaviour and density populations of the ragworm *Hediste diversicolor*?. *Marine Pollution Bulletin*, 2019, 142, pp.178-182. 10.1016/j.marpolbul.2019.03.026 . hal-02510702

HAL Id: hal-02510702

<https://univ-angers.hal.science/hal-02510702>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Is there a link between acetylcholinesterase, behaviour and density populations of the ragworm *Hediste diversicolor*?

Isabelle Métais^{a*}, Amélie Châtel^a, Mohammed Mouloud^a, Hanane Perrein-Ettajani^a,
Mélodie Bruneau^a, Patrick Gillet^a, Nisrine Jrad^b, Catherine Mouneyrac^a

^a Laboratoire Mer, Molécules, Santé (MMS, EA 2160), Université Catholique de l'Ouest, Faculté des Sciences, Angers F-49000 France

^b Laboratoire Angevin de Recherche en Ingénierie des Systèmes (LARIS, EA731), Université Catholique de l'Ouest, Faculté des Sciences, Angers F-49000 France

* Corresponding author: imetais@uco.fr

Abstract

The main objective of the present study was to explore the potential link between acetylcholinesterase (AChE) activity and burrowing behaviour of the ragworm *Hediste diversicolor*, which may have consequences at higher levels of biological organisation. Two complementary studies were conducted. AChE activity, at the sub-individual level, and behavioural responses, at the individual level, were evaluated in worms from the Loire estuary (France), whereas density and biomass of *H. diversicolor* were determined at the population level. A Spearman positive correlation between both biomarkers (AChE and burrowing) suggested that inhibition of AChE activity was linked to behaviour impairments. At the population level, lower AChE and behaviour activities were detected in worms corresponding to lower population density and biomass. These results provide direct empirical field evidence demonstrating the sensitivity of behaviour of *H. diversicolor* as a biomonitor of estuarine health status assessment.

Key words: *Hediste diversicolor*; acetylcholinesterase; population density; behaviour; Adverse Outcome Pathway.

Estuaries, as transitional zones, provide a wide range of vital nesting and feeding habitats and ecosystem services (Dallas and Jha, 2015), but since many decades, they are affected by multiple anthropogenic activities, leading to the release of various types of contaminants. To estimate the health status of ecosystems, biomonitoring programs aimed to classically combine chemical and biological approaches (e.g. multibiomarker approach) and ecological indicators. Conventional monitoring techniques are mainly based on the determination of the contaminants' concentrations in sediments and water. Evaluation of chemical contamination is a complex task, as estuaries often present a mixture of contaminants and their metabolites, some of which are costly or not yet accessible to analyze. It is widely recognised that chemical data alone can only provide partial information; for example, the unpredictability of the processes affecting the speciation of pollutants may alter their bioavailability and therefore their toxicity (Birch, 2018). For this reason, the integration of biomarker responses with chemical and ecological approaches on sentinel species allows evaluation of the toxicity of mixtures of pollutants and sub-lethal effects even at low concentrations (Amiard-Triquet et al., 2015). Biomarkers have been widely used to assess contaminant exposure and its effects in marine species such as the ragworm *Hediste diversicolor*, which plays an important role in the structure and functioning of marine ecosystems. However, biomarkers are subjected to the influence biotic and abiotic confounding factors (e.g. temperature, salinity, size/weight and reproductive status). Thus, to use these biomarkers in biomonitoring programs, it is necessary to previously determine baseline levels in species of interest, which is currently

challenging. Recently, Barrick et al. (2016; 2018) determined baseline assessment criteria (BAC) and environmental assessment criteria (EAC) of biochemical biomarkers (energy reserves, catalase: CAT, glutathione S transferases: GST, acetylcholinesterase: AChE and thiobarbituric acid reactive substances: TBARs) in *H. diversicolor* from a reference site (Durou et al., 2007). Regarding burrowing behaviour, Kalman et al. (2010) showed no influence of confounding factors (weight, salinity and grain size) on burrowing behaviour in *H. diversicolor*.

Field studies aiming at establishing a link between biomarkers at a low level of organisation and possible effects at a higher level of organisation (e.g. population and community) remain to be conducted. The adverse outcome pathway (AOP) approach proposed by Ankley et al. (2010) aims to break through the gap linking direct molecular initiating events with adverse outcomes at a higher level of organisation. AChE, a core biomarker of neurotoxicity, is well recognised as a biomarker of ecological relevance, as its activity inhibition observed in laboratory or field studies may lead to physiological and behavioural impairments in organisms exposed to not only pesticides but also metals, polycyclic aromatic hydrocarbons (PAHs), pharmaceutical products or algal toxins (Lionetto et al., 2013; Sturve et al., 2016; Lehtonen et al., 2003; Kankaanpää et al., 2007; Zhang et al., 2016).

Currently, the most robust of the developing pathway is the AOP linking AChE inhibition to acute mortality (Russom et al., 2014). This AOP has been proposed to support integrated approaches for testing and assessment (<https://aopwiki.org/aops/16>). However, there are many biochemical, behavioural, and physiological processes associated with the link between AChE inhibition and acute mortality of organisms and, consequently, possible density depletion of the species considered. The Loire estuary is a multi-contaminated estuary displaying a diffusive

pollution due to heavy metals, pesticides, PAHs and PCBs (Blanchet-Letrouvé et al., 2013), and AChE inhibition in organisms has been often linked to the presence of these contaminants. The main objective of the present study was to explore in the ragworm *H. diversicolor* from the Loire estuary (France), the potential link between AChE activity and burrowing behaviour that may have consequences at higher levels of biological organisation.

The sampling site was located on the largest mudflat of the polyhaline domain of the Loire estuary ('Les Brillantes') on the south shore downstream Paimboeuf (Fig. 1). Two zones were selected: the first one (Site 1: 47°16'55.70"N; 2° 3'47.60"W) was located in the upper intertidal area of the mudflat, 20 m from the shore, located close to an active eroded cliff (Thibault de Chanvalon et al., 2016). The second one, (Site 2: 47°17'3.77"N; 2° 3'42.87"W) was located in the middle intertidal area. Two sampling campaigns were conducted in May 2013 (after the flood period) and February 2014 (during the flood period), respectively.

For AChE activity determinations, 10 worms (*H. diversicolor*) in the same range of wet weight (mean S1: 0.220±0.114 g; mean S2: 0.187±0.121 g) were gently collected by hand, kept in wet sediment from their site of origin, immediately frozen at the laboratory and then kept at -80°C until analysis. AChE activity was determined using the method described in Buffet et al. (2011). Briefly, whole worm tissues were individually homogenised in 80 mM Tris-HCl pH 7.2, 200 mM NaCl, 1 mM Dithiothreitol supplemented with protease inhibitor cocktail according to the manufacturer's instructions. The homogenates were centrifuged at 9000 *g* for 30 min at 4°C, and supernatants were collected for protein (Bradford, 1976) and AChE activity determinations. Ten microlitres of the supernatant was added in triplicate to 190 µl 10 mM DTNB and 100 µL 100 mM acetylthiocholine into a 96-well microplate. The

99 reactions were read at 412 nm for 3 minutes. Results were expressed as nmoles of
100 thiocholine produced per min and per mg protein (nmoles min⁻¹ mg⁻¹ protein).
101 Concerning behavioural experiments, burrowing tests were set up in a dark
102 temperature-controlled room (15°C at the time of collection). Worms (*H. diversicolor*)
103 were submitted to burrowing tests as previously described by Bonnard et al. (2009).
104 Sediments from each site were homogenised by hand one day before experimentation.
105 Briefly, burrowing behaviour was studied by placing individually 20 worms on
106 sediments from their site of origin and recording their positions every 2 min for 30 min.
107 Automated behavioural tests were carried out with the Multifreshwater Biomonitor®
108 (MFB) from LimCo International according to the methodology previously described
109 and validated by Buffet et al. (2013). *H. diversicolor* movements inducing signals within
110 the range of 0.5 Hz were identified as undulation, whereas signals induced between
111 1.0 and 2.0 Hz indicated head movements and feeding activity. One individual per
112 chamber (n=7) was placed on sediment surface and allowed to acclimate 1 h before
113 MFB recording during 7 h. One control chamber without any organisms was used for
114 measuring background noise signals. Concerning density measurements, four
115 sediment samples from both areas (Site 1 and Site 2) were randomly collected at low
116 tide using plastic cores (10 cm x 35 cm) at a distance of 10 m from each other.
117 Sediment was sieved through a 1 mm mesh to collect and count worms. Fresh weight
118 of *H. diversicolor* individuals was determined to establish biomass. An average of 4
119 biomass replicates was calculated as the sum of fresh weight (g FW) divided by the
120 sediment surface sampled (m²) and expressed as FWg m⁻². Statistical analysis for
121 burrowing kinetics was performed on R (R Core Team 2014) using a generalised non-
122 linear model with α as burrowing rate (h⁻¹). This generalised non-linear model with a
123 conditional binomial stochastic part was fitted to each data set by using Bayesian

inference as described in Buffet et al. (2014). For all the other parameters, results are presented as mean \pm standard deviation, and statistical analyses were performed using XLSTAT 2012. Normality was assessed using the Shapiro–Wilk test. To define significant differences between sites for each sampling date, normally distributed data were analyzed by one-way ANOVA (multiple comparisons of mean values) followed by Tukey’s HSD post hoc method. When normality was not established, the non-parametric Kruskal–Wallis and Mann–Whitney comparison tests were used. The level of significance was established at $p < 0.05$. A Spearman's correlation (Daniel, 1990) was computed to quantify the relationship between AChE activity and behavioural toxicity parameter values. The Spearman's rank-order correlation is a non-parametric correlation coefficient ρ , measuring strength and direction of the monotonic relationship between AChE activity and MFB. The statistical significance test of the Spearman correlation was performed to investigate whether the null hypothesis ‘H0: There is no monotonic association between AChE and the behavioural toxicity parameter MFB’ can be rejected.

Lower AChE activity was found in worms (*H. diversicolor*) sampled from Site 2 (10.07 ± 5.80 nmole.min⁻¹.mg⁻¹ proteins) than those in Site 1 (23.34 ± 8.90 nmole.min⁻¹.mg⁻¹ proteins). AChE activities measured in *H. diversicolor* from Site 2 (compared to Site 1) showed 40% and 79.5% of inhibition in May 2013 and February 2014, respectively (Table 1). Results showing the positions of AChE activity measurements in worms from both areas (Site 1 and Site 2) according to the four colour traffic light previously identified by Barrick et al. (2018) (red: lethal effects expected; orange: sublethal effects expected; yellow: range expected for non-impacted site; green: evidence of good health) are illustrated in Figure 2. In February, AChE values were in the range where sub-lethal effects (worms from Site 1) and lethal effects (worms from

Site 2) were expected. In May, AChE values for worms from Site 1 were in the yellow range expected for non-impacted site, whereas those for worms from Site 2 were in the range where sub-lethal effects were expected. Results of burrowing kinetics carried out with *H. diversicolor* from both areas (Sites 1 and 2), allowed to burrow in sediment from their site of origin, are shown in Table 2. Burrowing kinetics of worms from Site 2 were significantly impaired in February 2014 compared to those in Site 1. Behavioural differences between worms from both sites measuring with the MFB are illustrated in Table 2. Significant frequency decrease ($p=0.021$) corresponding to body undulation (0.5 Hz) was observed for worms from Site 2 compared to those from Site 1. Decreases of head movements and feeding activity (between 1.0 and 2.0 Hz) were also observed for worms from Site 2 vs those from Site 1 ($p=0.066$). The mean Spearman's correlation coefficient of 0.5404 revealed a positive correlation ($p=0.0686$) between AChE activities and MFB values. Data of densities and biomass of *H. diversicolor* at both sites are shown in Table 3. Densities and biomass of ragworms collected at Site 1 were significantly higher than those at Site 2 for both sampling dates (May 2013 and February 2014).

The AOP approach initially proposed by Ankley et al. (2010) aims to link direct molecular initiating events with adverse outcomes at a higher biological level of organisation relevant to risk assessment. The usefulness of this approach for biomarker-based environmental risk assessments has been recently demonstrated using aquatic organisms (Wuk Lee et al., 2015). Effects observed at the individual and subcellular level (defence and damage responses: oxidative stress, immunotoxicity, neurotoxicity, genotoxicity and apoptosis) can be propagated upwards through increasing levels of organisation. Behavioural damages could find their origin in different biochemical and physiological impairments such as sensory system

174 disruption, endocrine disruption, neurotoxicity or energy disturbances (Amiard-Triquet
175 and Amiard, 2012) at a lower level and population effects at a higher level of
176 organisation. In the laboratory, numerous studies showed that decreased in AChE
177 activity induced behavioural disturbances in different species exposed to several
178 contaminants (Sismeiro-Vivas et al., 2007; Ren et al., 2015; Haverroth et al., 2015;
179 Geraudie et al., 2016). However, in the field, it is necessary to conduct studies able to
180 demonstrate a causal relationship between behavioural responses and effects at
181 higher levels of biological organisation such as population effects (Amiard-Triquet et
182 al., 2015). In the present study, we investigated, the potential links between three
183 different levels of biological organisation studied, precisely: sub-individual
184 (acetylcholinesterase activity), individual (burrowing behaviour) and population
185 (density and biomass) of worms. Results showed, at the biochemical level, a significant
186 lower value of AChE activity in worms (*H. diversicolor*) from Site 2 (decrease of 40%
187 in May 2013 and of 79% in February 2014 vs Site 1). Previously published studies
188 reported that a reduction of 20% in AChE activity in fish and invertebrates indicates
189 exposure to neurotoxic compounds (organophosphorus, pesticides), whereas
190 depletion of AChE activity by 20-50% indicates sublethal impact (Amiard-Triquet et al.,
191 2015). Mudflat sediments act as the final sink for various contaminants depending on
192 their properties. Sorption of organic contaminants on sediments is controlled by grain
193 size and organic matter content (Arslan-Alaton and Olmez-Hanci 2013). It has been
194 demonstrated that bioavailable metals and organic contaminant contents were low in
195 organic carbon enriched sites, tending to be adsorbed on fine-grained sediments
196 (Pelletier et al., 2011; Morrison et al., 2011). Site 1, compared to site 2, was composed
197 of slightly fine sediment and was slightly enriched in total organic carbon known to
198 have strong sorption capacities of contaminants (Luthy, 2004; Thibault de Chanvalon

et al., 2016). Moreover, Site 1 was located in an active eroded cliff where vegetation and rhizosphere could develop synergies with microorganisms increasing pollutant removal rates such as PAHs (Thibault de Chanvalon et al., 2016). At the individual level, burrowing kinetics of worms from Site 2 were significantly impaired compared to that in worms from Site 1. Schriks et al. (2006) used the MFB in *Xenopus laevis* as a sensitive and useful tool to quantify behavioural effects induced by toxic compound. In *H. diversicolor*, the use of MFB allows a more sensitive estimation of behavioural activities of worms, as differentiation between body undulation, head movements and feeding activity of worms can be assessed (Buffet et al., 2013). In this work, results showed lower body undulation of worms from Site 2 (vs Site 1). Statistical analysis (Spearman correlation) performed on the set results of AChE activities and behaviour (MFB) showing a positive correlation between both biomarkers (biochemical and behavioural) suggested that lower AChE activities were associated with lower behaviour of worms (and inversely). Feeding behaviour disturbances have been previously shown in *H. diversicolor* from the multipolluted Loire estuary as compared to controls from a reference site (Fossi-Tankoua et al., 2012). Recently, Harayashiki et al. (2016) demonstrated behavioural and AChE biochemical alterations in *Penaeus monodon* post larvae diet-exposed to inorganic mercury, reducing swimming activity. In the invertebrate *Gammarus fossarum*, Xuereb et al. (2009) showed that AChE inhibition levels greater than 50% led to population depletion. At the population level, worms exhibiting lower AChE activities and behaviour corresponded to those collected from the site (Site 2) where lower population densities were found. Indeed, densities of *H. diversicolor* showed a significantly contrasting status between both sites with significantly higher densities (280.61 ± 325.69 to 634.57 ± 294.45 ind.m⁻²) in the upper intertidal Site 1 and lower densities in the middle intertidal Site 2 (25.51 ± 23.29 to

66.96±28.28 ind.m⁻²). These observations cannot be relied on a patchy distribution of worms on the mudflat, as the distribution of *H. diversicolor* is known to be rather relatively uniform regarding the regularly distributed openings of burrows on the sediment surface (Kristensen et al., 2013). However, it has been demonstrated in *H. diversicolor* a double-way relationship between reworking behaviour and density as a decrease in individual sediment reworking activity when density increase (Duport et al. 2006) and at high levels of algal enrichment (Murray et al. 2017). Moreover, at high abundances, it has been shown that territorialism by this ragworm species is a mechanism underplaying trophic competition, while both phenomena seem to disappear at low population density (Kristensen et al. 2013). Site 1 was also enriched in living benthic foraminifera with a total density of 2199.35 individuals per 100 cm⁻³ compared to that at site 2, where a density of only 48 individuals per 100 cm⁻³ was found (Mojtahid et al., 2016). These authors correlated the microphytobenthic biofilm development on 'Les Brillantes' mudflat with the foraminifera species abundance, as it constitutes important quantities of food resources. Cardoso et al. (2004) showed that *H. diversicolor* had greater abundance in areas where algae were available, as a food source and a refuge from predators. Christensen et al. (2000) showed that an abundant supply of nutritious food (phytoplankton) increased *H. diversicolor* metabolism considerably. To cope, chemical stress is metabolically costly leading to a shift in allocation of energy with implications for growth and reproduction processes (Calow and Forbes, 1998). Thus, in the present study, it could be suggested that lower food resources and energetic cost of defence mechanisms to cope with chemical stress in worms from Site 2 vs Site 1 could lead to a reduction of the most energetic expensive individual processes, growth and reproductive potential, resulting in detrimental effects observed at the population level.

In conclusion, the contrasted status of *H. diversicolor* densities between both sites could be partly explained by AChE activity decrease and behavioural impairments of worms from the middle intertidal mudflat (Site 2) compared with the upper intertidal site (Site 1), which could result in the detrimental population effects observed. This study brings direct empirical field evidence supporting impact of AChE inhibition and behaviour at the population level demonstrating the sensitivity of behaviour as a biomonitor of estuarine health status assessment, as previously suggested by Roast et al. (2001). This integrated methodology should be included in biomonitoring surveys to evaluate habitat vulnerability and potential risk on higher trophic levels.

Acknowledgment. The research leading to these results has received funding from the RS2E-OSUNA project supported by the Région Pays de la Loire, France.

References

Amiard-Triquet, C., Amiard, J.C., 2012. Behavioral ecotoxicology. In: Amiard-Triquet, C., Amiard, J.C., Rainbow, P.S. (eds) Ecological biomarkers: indicators of ecotoxicological effects. 10, 253-277.

Amiard-Triquet, C., Amiard, J.C., Mouneyrac, C., 2015. Predictive ecotoxicology and environmental assessment. In: Amiard-Triquet, C., Amiard, J.C., Mouneyrac, C. (eds) Aquatic Ecotoxicology: advancing tools for dealing with emerging risks. chap19, 463-496.

Ankley, G.T., Bennett, R.S., Erickson, R.J., Hoff, D.J., Hornung, M.W., Johnson, R.D., Mount, D.R., Nichols, J.W., Russom, C.L., Schmieder, P.K., Serrano, J.A., Tietge, J.E.,

272 Arslan-Alaton, I., Olmez-Hanci, T., 2013. Sources of Environmental Pollution:
 273 Persistent Organic Pollutants. In: Suib, S.L. (ed) New and future developments in
 274 catalysis. 12, 303-398.

275 Barrick, A., Châtel, A., Marion, J.M., Perrein, H., Bruneau, M., Mouneyrac, C., 2016. A
 276 novel methodology for the determination of biomarker baseline levels in the marine
 277 polychaete *Hediste diversicolor*. Mar. Pollut. Bull. 108.
 278 10.1016/j.marpolbul.2016.04.056.

279 Barrick, A., Marion, J.M., Perrein, H., Châtel, A., Mouneyrac, C., 2018. Baseline levels
 280 of biochemical biomarkers in the endobenthic ragworm *Hediste diversicolor* as useful
 281 tools in biological monitoring of estuaries under anthropogenic pressure. Mar. Pollut.
 282 Bull. 129, 81-85. 10.1016/j.marpolbul.2018.02.006.

283 Benyoucef, I., Blandin, E., Lerouxel, A., Jesus, B., Rosa, P., Méléder, V., Launeau, P.,
 284 Barillé, L., 2014. Microphytobenthos interannual variations in a north-European
 285 estuary (Loire estuary, France) detected by visible-infrared multispectral remote
 286 sensing. Estuar. Coast. Shelf Sci. 136, 43-52.

287 Birch, G.F., 2018. A review of chemical-based sediment quality assessment
 288 methodologies for the marine environment. Mar. Pollut. Bull. 133, 218-232.

289 Blanchet-Letrouvé, I., Lafont, A.G., Poirier, L., Baloché, S., Zalouck-Vergnoux, A.,
 290 Dufour, S., Mouneyrac, C., 2013. Vg mRNA induction in an endangered fish species
 291 (*Anguilla anguilla*) from the Loire estuary (France). Ecotoxicol. Environ. Saf. 97, 103-
 292 113.

293 Bonnard, M., Roméo, M., Amiard-Triquet, C., 2009. Effects of copper on the burrowing
 294 behavior of estuarine and coastal invertebrates, the polychaete *Nereis diversicolor* and
 295 the bivalve *Scrobicularia plana*. Hum. Ecol. Risk Assess. 15, 11-26.

296 Bradford, M., 1976. A rapid and sensitive method for the quantitation of microgram
 297 quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72,
 298 248-254.

299 Buffet, P.E., Tankoua, O.F., Pan, J.F., Berhanu, D., Herrenknecht, C., Poirier, L.,
 300 Amiard-Triquet, C., Amiard, J.C., Bérard, J.B., Risso, C., Guibbolini, M., Roméo, M.,
 301 Reip, P., Valsami-Jones, E., Mouneyrac, C., 2011. Behavioural and biochemical
 302 responses of two marine invertebrates *Scrobicularia plana* and *Hediste diversicolor* to
 303 copper oxide nanoparticles. *Chemosphere.* 84(1), 166-74. doi:
 304 10.1016/j.chemosphere.2011.02.003.

305 Buffet, P.E., Poirier, L., Zalouk-Vergnoux, A., Lopes, C., Amiard, J.C., Gaudin, P.,
 306 Risso-de Faverney, C., Guibbolini, M., Gilliland, D., Perrein-Ettajani, H., Valsami-
 307 Jones, E., Mouneyrac, C., 2014. Biochemical and behavioural responses of the marine
 308 polychaete *Hediste diversicolor* to cadmium sulfide quantum dots (CdS QDs):
 309 Waterborne and dietary exposure. *Chemosphere* 100, 53-70.

310 Buffet, P.E., Richard, M, Caupos, F., Vergnoux, A., Perrein-Ettajani, H., LunaAcosta,
 311 A., Akcha, F., Amiard, J.C., Amiard-Triquet, C., Guibbolini, M., Risso-de Faverney, C.,
 312 Thomas-Guyon, H., Reip, P., Dybowska, A., Berhanu, D., Valsami-Jones, E.,
 313 Mouneyrac, C., 2013. Mesocosm study of fate and effects of CuO nanoparticles on
 314 endobenthic species (*Scrobicularia plana*, *Hediste diversicolor*). *Environ. Sci. Technol.*
 315 47, 1220-1228.

316 Calow, P., Forbes, V.E., 1998. How do physiological responses to stress translate into
 317 ecological and evolutionary processes? *Comp. Biochem. Physiol. A* 120, 11–16.

318 Cardoso, P.G., Pardal, M.A., Raffaelli, D., Baeta, A., Marques, J.C., 2004.
 319 Macroinvertebrate response to different species of macroalgal mats and the role of
 320 disturbance history. *J. Exp. Mar. Biol. Ecol.* 308, 207-220.

321 Christensen, B., Vedel, A., Kristensen, E., 2000. Carbon and nitrogen fluxes in
 322 sediment inhabited by suspension-feeding (*Nereis diversicolor*) and non-suspension-
 323 feeding (*N. virens*) polychaetes. Mar. Ecol. Prog. Ser. 192, 203-217.

324 Dallas, L.J., Jha, A.N., 2015. Applications of biological tools or biomarkers in aquatic
 325 biota: A case study of the Tamar estuary, South West England. Mar. Pollut. Bull. 95,
 326 618-633.

327 Daniel, W.W., 1990. Spearman Rank Correlation Coefficient. Applied Nonparametric
 328 Statistics (2nd ed., pp. 358-365). Boston, MA: PWS-Kent.

329 Duport, E., Stora, G., Tremblay, P., Gilbert, F., 2006. Effects of population density on
 330 the sediment mixing induced by the gallery-diffusor *Hediste (Nereis) diversicolor* O.F.
 331 Muller, 1776. J. Exp. Mar. Biol. Ecol. 336, 33-41.

332 Durou, C., Poirier, L., Amiard, J.C., Budzinski, H., Gnassia-Barelli, M., Lemenach, K.,
 333 Peluhet, L., Mouneyrac, C., Roméo, M., Amiard-Triquet, C., 2007. Biomonitoring in a
 334 clean and multi-contaminated estuary based on biomarkers and chemical analyses in
 335 the endobenthic worm *Nereis diversicolor*. Environ. Pollut. 148, 445-458.

336 Fossi Tankoua, O., Buffet, P.E., Amiard, J.C., Amiard-Triquet, C., Méléder, V., Gillet,
 337 P., Mouneyrac, C., Berthet, B., 2012. Intersite variations of a battery of biomarkers at
 338 different levels of biological organisation in the estuarine endobenthic worm *Nereis*
 339 *diversicolor* (Polychaeta, Nereididae) Aquat. Toxicol. 114-115, 96-103.

340 Geraudie, P., Bakkemo, R., Milinkovitch, T., Thomas-Guyon, H., 2016. First evidence
 341 of marine diesel effects on biomarker responses in the Icelandic scallops, *Chlamys*
 342 *islandica*. Environ. Sci. Pollut. Res. Int. 23, 16, 16504-16512.

343 Harayashiki, C.A.Y., Reichelt-Brushett, A.J., Liu, L., Butcher, P., 2016. Behavioural
 344 and biochemical alterations in *Penaeus monodon* postlarvae diet-exposed to inorganic
 345 mercury. Chemosphere 164, 241-247.

346 Haverroth, G.M., Welang, C., Mocelin, R.N., Postay, D., Bertoncello, K.T.,
 347 Franscescon, F., Rosemberg, D.B., Dal Magro, J., Dalla Corte, C.L., 2015. Copper
 348 acutely impairs behavioral function and muscle acetylcholinesterase activity in
 349 zebrafish (*Danio rerio*). *Ecotoxicol. Environ. Saf.* 122, 440-447.

350 Kalman, J., Buffet, P.E., Amiard, J.C., Denis, F., Mouneyrac, C., Amiard-Triquet, C.,
 351 2010. Assessment of the influence of confounding factors (weight, salinity) on the
 352 response of biomarkers in the estuarine polychaete *Nereis diversicolor*. *Biomarkers*
 353 15, 5, 461-469.

354 Kankaanpää, H., Leiniö, S., Olin, M., Sjövall, O., Meriluoto, J., Lehtonen, K.K., 2007.
 355 Accumulation and depuration of cyanobacterial toxin nodularin and biomarker
 356 responses in the mussel *Mytilus edulis*. *Chemosphere* 68, 7, 1210-1217.

357 Kristensen, E., Delefosse, M., Quintana, C.O., Banta, G.T., Petersen, H.C., Jorgensen,
 358 B., 2013. Distribution pattern of benthic invertebrates in Danish estuaries: The use of
 359 Taylor's power law as a species-specific indicator of dispersion and behavior. *J. Sea*
 360 *Res.* 77, 70-78.

361 Lehtonen, K.K., Kankaanpää, H., Leiniö, S., Sipiä, V.O., Pflugmacher, S., Sandberg-
 362 Kilpi, E., 2003. Accumulation of nodularin-like compounds from the cyanobacterium
 363 *Nodularia spumigena* and changes in acetylcholinesterase activity in the clam *Macoma*
 364 *balthica* during short-term laboratory exposure. *Aquat. Toxicol.* 64, 461-476.

365 Lionetto, M.G., Caricato, R., Calisi, A., Giordano, M.E., Schettino, T., 2013.
 366 Acetylcholinesterase as a biomarker in environmental and occupational medicine: New
 367 insights and future perspectives. *Biomed. Res. Int.* ID 321213
 368 <http://dx.doi.org/10.1155/2013/321213>.

369 Luthy, R.G., 2004. Organic contaminants in the environment: challenges for the
 370 water/environmental engineering community. In: Norling, P., Wood-Black, F.,

371 Masciangioli, T.M. (eds). Water and Sustainable Development: Opportunities for the
 372 Chemical Sciences: A Workshop Report to the Chemical Sciences Roundtable
 373 Washington (DC): National Academies Press (US).
 374 Mojtahid, M., Geslin, E., Coynel, A., Gorse, L., Vella, C., Davranche, A., Zozzolo, L.,
 375 Blanchet, L., Bénéteau, E., Maillet, G., 2016. Spatial distribution of living (Rose Bengal
 376 stained) benthic foraminifera in the Loire estuary (western France). J. Sea Res. 118,
 377 1-16.
 378 Morrison, G., Greening, H., Yates, K.K., 2011. Sediment contaminants and benthic
 379 habitat quality. In: Morrison G, Greening H, Yates KK (eds) Integrating Science and
 380 Resource Management in Tampa Bay, Florida. US Geological Survey Circular 1348,
 381 7, 203-238.
 382 Murray, F., Solan, M., Douglas, A., 2017. Effects of algal enrichment and salinity on
 383 sediment particle reworking activity and associated nutrient generation mediated by
 384 the intertidal polychaete *Hediste diversicolor*. J. Exp. Mar. Biol. Ecol. 495, 75-82.
 385 Pelletier, M.C., Campbell, D.E., Ho, K.T., Burgess, R.M., Audette, C.T., Detenbeck,
 386 N.E., 2011. Can sediment total organic carbon and grain size be used to diagnose
 387 organic enrichment in estuaries? Environ. Toxicol. Chem. 30, 3, 538-547.
 388 Ren, Z., Zhang, X., Wang, X., Qi, P., Zhang, B., Zeng, Y., Fu, R., Miao, M., 2015. AChE
 389 inhibition: One dominant factor for swimming behavior changes of *Daphnia magna*
 390 under DDVP exposure. Chemosphere. 120: 252-257.
 391 Roast, S.D., Widdows, J., Jones, M.B., 2001. Impairment of mysid (*Neomysis integer*)
 392 swimming ability: an environmentally realistic assessment of the impact of cadmium
 393 exposure. Aquat. Toxicol. 52, 217-227.

394 Russom, C.L., Lalone, C.A., Villeneuve, D.L., Ankley, G.T., 2014. Development of an
 395 adverse outcome pathway for acetylcholinesterase inhibition leading to acute mortality.
 396 Environ. Toxicol. Chem. 33, 10, 2157-2169.

397 Schriks, M., van Hoorn, M.K., Faassen, E.J., van Dam, J.W., Murk, A.J., 2006. Real-
 398 time automated measurement of *Xenopus leavis* tadpole behavior and behavioral
 399 responses following triphenyltin exposure using the multispecies freshwater biomonitor
 400 (MFB). Aquat. Toxicol. 77, 298-305.

401 Sismeiro-Vivas, J., Abrantes, N., Pereira, J.L., Castro, B.B., Gonçalves, F., 2007.
 402 Short-term effects of Quirlan (chlorfenvinphos) on the behavior and
 403 acetylcholinesterase activity of *Gambusia holbrooki*. Environ Toxicol. 22(2), 194-202.

404 Sturve, J., Scarlet, P., Halling, M., Kreuge, J., Macia, A., 2016. Environmental
 405 monitoring of pesticide exposure and effects on mangrove aquatic organisms of
 406 Mozambique. Mar. Environ. Res. 121, 9-19.

407 Thibault de Chanvalon, A., Mouret, A., Knoery, J., Geslin, E., Péron, O., Metzger, E.,
 408 2016. Manganese, iron and phosphorus cycling in an estuarine mudflat sediment,
 409 Loire, France. J. Sea Res. 118, 92-102.

410 Wuk Lee, J., Won, E.J., Raisuddin, S., Lee, J.S., 2015. Significance of adverse
 411 outcome pathways in biomarker-based environmental risk assessment in aquatic
 412 organisms. J. Environ. Sci. 35, 115-127.

413 Xuereb, B., Lefèvre, E., Garric, J., Geffard, O., 2009. Acetylcholinesterase activity in
 414 *Gammarus fossarum* (Crustacea Amphipoda): Linking AChE inhibition and
 415 behavioural alteration. Aquat. Toxicol. 94, 114-122.

416 Zhang, D.L., Liu, S.Y., Zhang, J., Zhang, J.K., Hu, C.X., Liu, Y.D., 2016. Respiratory
 417 toxicity of cyanobacterial aphantoxins from *Aphanizomenon flos-aquae* DC-1 in the
 418 zebrafish gill. Aquat. Toxicol. 176, 106-115.




Fig. 1. Brillantes mudflat is localized in the polyhaline domain of the Loire estuary (France), with its intertidal flats shown in dark gray (Benyoucef et al., 2014).

Table 1: AChE activity measured in *H. diversicolor* from site 1 and site 2 expressed in nmole.min⁻¹.mg⁻¹ proteins. ^a significantly difference compared to ^b (p<0.05).

	AChE activity (nmole.min ⁻¹ .mg ⁻¹ proteins) Site1	AChE activity (nmole.min ⁻¹ .mg ⁻¹ proteins) Site2	P value
May 2013	19.75±5.01 ^a	11.83±5.70 ^b	0.008
February 2014	30.64±12.42 ^a	6.54±4.61 ^b	0.002




Fig. 2 BAC (baseline assessment criteria) and EAC (environmental assessment criteria) graph for AChE activity values (Barrick et al., 2018). The black line indicates the mean values established for the low-impact site Authie, the yellow band indicates the range of the BAC and the orange line indicates the range where sublethal effects are expected to occur. AChE values for both sites (Site 1: S1; Site 2: S2) on May 2013 (May 13) and February 2014 (Feb 14) were plotted on the graph as well.

Table 2: Burrowing tests: Mean estimate α (h^{-1}) and the 95% credibility interval (CI) obtained after *a posteriori* distribution for each site (site 1: S1, site 2: S2) and sampling period (May 2013 and February 2014). MFB: Mean frequency occurrence (%) and standard deviation (SD) of signals recorded with Multi Freshwater Biomonitor (MFB) and corresponding to body undulation (0.5 Hz) and to head movements and feeding

445 activity (1 to 2 Hz) in *H. diversicolor*. * significant difference ($p < 0.05$). nd: not
 446 determined.

	Burrowing test α [CI] S1	Burrowing test α [CI] S2	MFB Frequency occurrence (mean% \pm SD) 0.5 Hz S1	MFB Frequency occurrence (mean% \pm SD) 0.5 Hz S2	MFB Frequency occurrence (mean% \pm SD) 1.2 Hz S1	MFB Frequency occurrence (mean% \pm SD) 1.2 Hz S2
May 2013	0.064 [0.036 - 0.099]	0.059 [0.033 - 0.091]	64.89 \pm 9.13*	43.95 \pm 16.15 * ($p=0.021$)	10.48 \pm 5.88	7.27 \pm 2.01 ($p=0.523$)
February 2014	0.103 [0.063 - 0.152]*	0.036 [0.020 - 0.059]*	58.00 \pm 4.43	49.51 \pm 7.70 ($p=0.066$)	6.44 \pm 0.97	5.61 \pm 0.92 ($p=0.173$)

447

448 Table 3. Intersite variations of *H. diversicolor* density (number of individuals N m⁻² as
 449 mean \pm standard deviation) and biomass (Fresh weight FWg m⁻² as mean \pm standard
 450 deviation) for each sampling period (May 2013 and February 2014). nd: not determined

	Density (N.m ⁻²) Site 1	Density (N.m ⁻²) Site 2	Biomass (FWg.m ⁻²) Site 1	Biomass (FWg.m ⁻²) Site 2
May 2013	631.38 \pm 182.03	51.02 \pm 31.24	39.80 \pm 21.13	3.18 \pm 1.96
February 2014	280.61 \pm 325.69	66.96 \pm 28.28	19.84 \pm 23.26	4.46 \pm 4.64

451