



HAL
open science

Tetrathiafulvalene-[2.2]paracyclophanes: Synthesis, chiral structures and chiroptical properties

Cécile Mézière, Magali Allain, Cristina Oliveras-Gonzalez, Thomas Cauchy, Nicolas Vanthuyne, Laura Sarbu, Lucian Birsa, Flavia Pop, Narcis Avarvari

► **To cite this version:**

Cécile Mézière, Magali Allain, Cristina Oliveras-Gonzalez, Thomas Cauchy, Nicolas Vanthuyne, et al.. Tetrathiafulvalene-[2.2]paracyclophanes: Synthesis, chiral structures and chiroptical properties. Journées André Collet de la Chiralité, 2018, Noirmoutier, France. , 2018. hal-02564566

HAL Id: hal-02564566

<https://univ-angers.hal.science/hal-02564566v1>

Submitted on 5 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Cécile Mézière¹, Magali Allain¹, Cristina Oliveras-González¹, Thomas Cauchy¹, Nicolas Vanthuyne²,
Laura G. Sarbu³, Lucian M. Birsa³, Flavia Pop¹, Narcis Avarvari¹

¹Laboratoire MOLTECH-Anjou, UMR6200 CNRS-Université d'Angers, Faculté des Sciences, Bât. K, 2 Bd Lavoisier, 49045 Angers Cedex, France

²Aix Marseille Université, CNRS, Centrale Marseille, iSm2, Marseille, France


³Department of Chemistry, "Al. I. Cuza", University of Iasi, Iasi, Romania

Objective: Planar chirality is introduced in electroactive tetrathiafulvalenes via paracyclophanes units in order to explore the chiroptical properties of the new compounds formed. The challenge is also to investigate the influence of chirality on conductivity properties of the corresponding mixed valence salts that could be obtained by electrocrystallization as previously described in our group [2].



The two enantiomers of monosubstituted [2.2]paracyclophane: R_p S_p

Synthesis: heterocoupling in the presence of phosphite



Purification of the coupling products is achieved by chromatography on silica gel ; The bis [2.2]paracyclophane TTF is not formed.

Enantiomer resolution by chiral HPLC:

On Chiralpak IE (250 x 10 mm), hexane/ethanol/dichlorométhane (5/3/2), 5 mL/min, UV detection at 254nm.

(rac)-1 → (-)-enantiomer ee > 98.5% (S_p)
(rac)-1 → (+)-enantiomer ee > 99% (R_p)
(rac)-2 → (-)-enantiomer ee > 99%
(rac)-2 → (+)-enantiomer ee > 99%

Cyclic voltammetry:



Two one electron reversible oxidation processes for (rac) 1 and 2

Compound	E ¹ _{ox}	E ² _{ox}
1	0.02	0.47
2	0.14	0.62
TTF-(SMe) ₂ [3]	0.03	0.35
Ph-TTF-(SMe) ₂ [4]	0.00	0.37

No significant influence of the paracyclophane unit on the oxidation potentials.

The electron withdrawing effect of the carbonyl units in 2 reduces its donor ability.


Half wave potentials relative to Fc/Fc⁺, DCM 10⁻¹ M TBAFP₆, 20 ° C

Frontier molecular orbitals:

MO Energy

LUMO -1.42 eV


HOMO -4.94 eV



MO Energy

LUMO -1.97 eV

HOMO -5.11 eV




The HOMO corresponds to the TTF moiety and the stability is higher for compound 2 ; In 1 the LUMO is based on the cyclophane unit while in 2 it is spread over the ester part.

Crystal structures:


The crystals were obtained after slow evaporation of an acetonitrile solution in a very thin tube [4]

(R_p)-1

Crystal size: 0.216 x 0.148 x 0.038 mm
T = 200.0(1) K, λ = 1.54184 Å
a = 7.4706(2) Å α = 90°
b = 11.7648(3) Å β = 100.454(3)°
c = 13.0617(4) Å γ = 90°
V = 1128.94(5) Å³
R₁ = 4.80%
Monoclinic non centrosymmetric space group, P 2₁




Packing of (R_p)-1 in the ab plane



ArH...π interactions between the alternating stacks


Packing of (R_p)-1 in the bc plane




Lateral short S...S distances between the TTF units of the orthogonal molecules

(rac)-2

Crystal size: 0.394 x 0.030 x 0.023 mm
T = 150.0(1) K, λ = 1.54184 Å
a = 17.0529(9) Å α = 90°
b = 12.6806(3) Å β = 113.142(6)°
c = 17.1325(9) Å γ = 90°
V = 3406.6(3) Å³
R₁ = 8.31%
Monoclinic non centrosymmetric space group, P 2₁




Packing of (rac)-2 in the ab plane



Hydrogen bond interactions of the ester units


(rac)-2 ArH...π interactions between layers




Segregation of the TTF and the p-cyclophanes units

ArH...π interactions between the alternating stacks but no S...S one : high distortion compared to (R_p)-1 (less favorable packing)

Circular dichroism:



First intense band at 296 nm, two bands at 325 (S₀ to S₄ excitation) and 350 nm, followed by one large band of weak intensity for enantiomers of 2, and two bands of low intensity at 395 (S₀ to S₃ excitation) and 435 nm (S₀ to S₂ excitation) for the enantiomers of compound 1. There is a small blue shift of all bands compare to the calculated data. The TD-DFT calculation method is not suitable for compound 2 because of gas phase position of ester groups, heavily distorted from the TTF median plan, generating an artificial CD signal.



[1] Mézière C., Allain M., Oliveras-González C., Cauchy T., Vanthuyne N., Sarbu L.G., Birsa L.M., Pop F., *Chirality*, 2018, 30, 568.

[2] Pop F., Auban-Senzier P., Canadell E., Rikken G.L.J.A., Avarvari N., *Nat. Commun.*, 2014, 5, 3757.

[3] Sarbu L.G., Bahrin L.G., Jones P.G., Birsa L.M., Hopf H., *Beilst. J. Org. Chem.*, 2015, 11, 1917.

[4] Spingler B., Schnidrig S., Todorova T., Wild F., *Cryst. Eng. Comm.*, 2012, 14, 751.

[5] Kobayakawa K., Hasegawa M., Sasaki H., Endo J., Matsuzawa H., Sako K., Yoshida J., Mazaki Y., *Chem. Asian J.*, 2014, 9, 2751.