



**HAL**  
open science

## The dielectric constant and solvent effect on the stabilities of copper(II) chloro complexes.

Mustayeen A. Khan, Gilles M. Bouet, Marie-José Schwing.

► **To cite this version:**

Mustayeen A. Khan, Gilles M. Bouet, Marie-José Schwing.. The dielectric constant and solvent effect on the stabilities of copper(II) chloro complexes.. *Transition Metal Chemistry*, 1997, 22, pp.604-605. 10.1023/A:1018577127619 . hal-03193553

**HAL Id: hal-03193553**

**<https://univ-angers.hal.science/hal-03193553>**

Submitted on 11 Apr 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Communication

# The dielectric constant and solvent effect on the stabilities of copper(II) chloro complexes

Mustayeen A. Khan\* and Gilles M. Bouet

Laboratoire de Chimie de Coordination, Faculté de Pharmacie, 16 Boulevard Daviers, F-49100 Angers, France

Marie-José Schwing

Laboratoire de Chimie Physique, E.H.I.C.S., Rue Blaise Pascal, F-67000 Strasbourg, France

The individually published results of the stability of copper(II) chloro complexes in various solvents have been assembled. The presence of four mononuclear complexes was postulated in the following solvents: ethanoic acid, tetrahydrofuran, trimethylphosphate, acetone, methanol, propylene carbonate and water, whereas dimethylformamide and dimethyl sulfoxide are characterized by the presence of only three complexes. The respective stability constants,  $\beta_j$  have been calculated and it is shown here that, for the protic solvents,  $\log \beta_j$  is an inverse linear function of the dielectric constant,  $\epsilon_r$ , of the solvent. However, for aprotic solvents, stability decreases with the increasing value of  $\epsilon_r$  up to a certain value and then increases sharply from  $\epsilon_r \approx 45$ . This sudden increase in the stability of complexes in propylene carbonate is explained on the basis of its low value for the molecular weight to dielectric constant ratio ( $\mu_\epsilon$ ). This ratio should be given more importance in the interpretation of results correlating the stabilities with a physical parameter than the values of the dielectric constant alone, which are in themselves not static but a function of solution concentrations.

During the last fifteen years, a rigorous study of the copper(II) chloro complexes has been undertaken in different solvents. These studies determined the number of complexes formed in solution, their nature, their stability constants and their electronic spectra, and have been extended to other transition metals such as cobalt(II) and nickel(II)<sup>(1–3)</sup>.

For the cupric ion, Table 1 summarises the results of the equilibrium studied:  $\text{Cu}^{2+} + j\text{Cl}^- \rightleftharpoons \text{CuCl}_j^{(j-2)-}$  where  $j \leq 4$ . The equilibrium is defined by the overall stability constant  $\beta_j = [\text{CuCl}_j^{(j-2)-}] / [\text{Cu}^{2+}][\text{Cl}^-]^j$ .

The results were obtained under the same experimental conditions, with the same methodology and using the same calculation program. Since we are comparing results in different solvents, the bibliography is limited to work carried out under similar conditions.

The presence of four successive mononuclear complexes was postulated in all the solvents studied except DMF and DMSO, where the absence of the second complex,  $\text{CuCl}_2$ , is accounted for by autocomplex formation leading to  $\text{CuCl}^+$  and  $\text{CuCl}_3^-$ . The stability constants of all the species were derived using the same method of calculation and the solvent effect in certain categories discussed. A peculiar behaviour, however, is now reported and this, of course, became apparent only

when all the individually published results were assembled.

A plot of  $\log \beta_j$  versus the dielectric constant of the solvent,  $\epsilon_r$ , shows that for protic solvents AcOH, MeOH and H<sub>2</sub>O the stability diminishes with increasing value of the dielectric constant (Figure 1).  $\log \beta_j$  is therefore a linear function of  $\epsilon_r$ .

However, for the aprotic solvents studied: THF, TMP, Me<sub>2</sub>CO, DMF, DMSO and propylene carbonate (PC), it can be discerned clearly that stability decreases until a certain value of dielectric constant, *ca.* 45 is reached, and then it increases sharply. This critical dip is interesting and can be further confirmed by studies of this system in solvents with  $\epsilon_r$  values between 40 and 70.

This steep increase in the stability in PC has been reported by various authors even when the results were obtained by other experimental methods. Scharff<sup>(12)</sup>, using potentiometric methods, reported the presence of three complexes in this medium and calculated their stability constants as  $\log \beta_{\text{CuCl}^+} = 12.2$ ,  $\log \beta_{\text{CuCl}_2} = 20.8$  and  $\log \beta_{\text{CuCl}_3^-} = 27.4$ . One of the reasons for these high values in PC is probably the low value of the ratio of its molecular weight to the dielectric constant:  $\mu_\epsilon = \text{mw}/\epsilon_r = 102/69 = 1.48$ . For all other aprotic solvents  $\mu_\epsilon$  has a higher value, *viz.* DMSO = 1.74; DMF = 2.03; Me<sub>2</sub>CO = 2.77; TMP = 6.80 and THF = 9.49.

We believe that the ratio of the molecular weight to the dielectric constant should be given more importance *vis-à-vis* the dielectric constant of the solvent alone, which is not a static value for the purpose of generalisation. The molecular weight of the solvent, on the other hand, is an inherent and immutable property.

It may be recalled that the presence of ions in a solvent causes a decrease in the dielectric constant and therefore each solution containing a different ligand concentration has a different dielectric value. The reason for this decrease in the presence of ions is probably due to the restricted rotation of molecules in the vicinity of an ion, which hinders their orientation in the external field. The orientation of molecules can therefore no longer be controlled by the external field and the dielectric constant decreases with increasing concentration<sup>(13)</sup>.

However, attempts to plot  $\log \beta_j$  versus  $\mu_\epsilon$  yield a complex function, not at all like the simple linearity of  $\log \beta_j$  versus  $\epsilon_r$ . In conclusion, although studies on the basis of the dielectric constant of the solvent alone are simpler and give a fairly approximate idea of the stabilities of the complexes in different solvents, extreme care must be taken in generalising this behaviour.

\* Author to whom all correspondence should be directed.

Table 1. Logarithm of stability constants of copper(II) chloro complexes identified in different solvents

Solvent	$\epsilon_r^a$	$\log \beta_1$	$\log \beta_2$	$\log \beta_3$	$\log \beta_4$	Ref.
Ethanoic acid (AcOH)	6.2	3.4	5.6	7.4	7.8	4
Tetrahydrofuran (THF)	7.6	11.8	20.3	23.9	25.7	5
Trimethylphosphate (TMP)	20.6	7.3	12.4	16.8	19.4	6
Acetone (Me <sub>2</sub> CO)	21.0	8.4	14.5	19.9	21.5	7
Methanol (MeOH)	32.6	2.5	4.2	5.4	5.7	8
Dimethylformamide (DMF)	36.0	3.8	—	9.8	10.8	9
Dimethyl sulfoxide (DMSO)	45.0	3.0	—	6.9	7.5	10
Propylene carbonate (PC)	69.0	12.0	22.7	28.9	34.1	10
Water	78.5	0.60	0.67	0.29	-0.64	11

<sup>a</sup> Dielectric constant of the solvent.




Figure 1. A plot of  $\log \beta_j$  of the copper(II) chloro complexes versus the dielectric constant  $\epsilon_r$  of the nine solvents studied. Dotted lines, protic solvents; solid lines, aprotic solvents.

## References

- (1) S. Lechat, M. A. Khan, G. Bouet and F. Vierling, *Inorg. Chim. Acta*, **211**, 33 (1993).
- (2) M. A. Khan, D. Cronier, G. Boeut and F. Vierling, *Transition Met. Chem.*, **20**, 369 (1995).
- (3) M. A. Khan, G. Bouet, F. Vierling, J. Meullemeestre and M. J. Schwing, *Transition Met. Chem.*, **21**, 231 (1996).
- (4) M. A. Khan, J. Meullemeestre, M. J. Schwing and F. Vierling, *Polyhedron*, **2**, 459 (1983).
- (5) C. Amuli, J. Meullemeestre, M. J. Schwing and F. Vierling, *Inorg. Chem.*, **22**, 3567 (1983).
- (6) C. Amuli, M. Elleb, J. Meullemeestre, M. J. Schwing and F. Vierling, *Inorg. Chem.*, **25**, 856 (1986).
- (7) C. Amuli, J. Meullemeestre, M. J. Schwing and F. Vierling, *Nouv. J. Chim.*, **11**, 27 (1987).
- (8) M. A. Khan, J. Meullemeestre, M. J. Schwing and F. Vierling, *Inorg. Chem.*, **28**, 3306 (1989).
- (9) M. Elleb, J. Meullemeestre, M. J. Schwing and F. Vierling, *Inorg. Chem.*, **19**, 2699 (1980).
- (10) M. Elleb, J. Meullemeestre, M. J. Schwing and F. Vierling, *Inorg. Chem.*, **21**, 1477 (1982).
- (11) M. A. Khan and M. J. Schwing-Weill, *Inorg. Chem.*, **15**, 2202 (1976).
- (12) J. P. Scharff, *Bull. Soc. Chim. Fr.*, **1**, 413 (1972).
- (13) G. Kortüm, *Treatise on Electrochemistry*, Elsevier, Amsterdam, 1965, Ch. III.

(Received 18 June 1997)

Accepted 11 July 1997)

TMC 4004(C)