



Persistent organochlorine pesticides in periprostatic adipose tissue from men with prostate cancer: Ethno-geographic variations, association with disease aggressiveness

Jean-Philippe Antignac, Sandy Figiel, Michèle Pinault, Pascal Blanchet, Franck Bruyère, Romain Mathieu, Souhil Lebdai, Georges Fournier, Jérôme Rigaud, Karine Mahéo, et al.

► To cite this version:

Jean-Philippe Antignac, Sandy Figiel, Michèle Pinault, Pascal Blanchet, Franck Bruyère, et al.. Persistent organochlorine pesticides in periprostatic adipose tissue from men with prostate cancer: Ethno-geographic variations, association with disease aggressiveness. *Environmental Research*, 2023, 216 (Pt 4), pp.114809. 10.1016/j.envres.2022.114809 . hal-03898019

HAL Id: hal-03898019

<https://hal.science/hal-03898019>

Submitted on 25 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PERSISTENT ORGANOCHLORINE PESTICIDES IN PERIPROSTATIC ADIPOSE TISSUE FROM MEN WITH PROSTATE CANCER: ETHNO-GEOGRAPHIC VARIATIONS, ASSOCIATION WITH DISEASE AGGRESSIVENESS.

Jean-Philippe Antignac¹, PhD, Sandy Figiel², PhD, Michèle Pinault², MS, Pascal Blanchet^{3,4}, PhD, Franck Bruyère⁵, PhD, Romain Mathieu^{4,6}, PhD, Souhil Lebdaï⁷, PhD, Georges Fournier⁸, PhD, Jerome Rigaud⁹, PhD, Karine Mahéo², PhD, Philippe Marchand¹, Ingrid Guiffard¹, Emmanuelle Bichon¹, Bruno le Bizec¹, PhD, Luc Multigner⁴, PhD, and Gaëlle Fromont^{2,5}, PhD.

¹Oniris, INRAE, LABERCA, 44300 Nantes, France.

²Inserm UMR1069 “Nutrition, Croissance et Cancer” Université François Rabelais, Faculté de Médecine, 10 bd Tonnellé, 37032 Tours, France

³CHU Pointe à Pitre, Department of Urology

⁴Inserm UMR1085 - IRSET Rennes

⁵CHRU Bretonneau, Departments of Pathology and Urology, Tours, France

⁶CHU Rennes, Departments of Pathology and Urology

⁷CHU Angers, Department of Urology

⁸CHU Brest, Departments of Pathology and Urology

⁹CHU Nantes, Departments of Pathology and Urology

Corresponding author: Gaëlle FROMONT

Address: INSERM U1069, “Nutrition, Croissance et Cancer”, Université François-Rabelais, Faculté de Médecine, 10 Bd Tonnellé, F-37032 TOURS Cedex, France. Tél: (+33) 2 47 36 62 13. Fax : (+33) 2 47 36 62 26.

E-mail address: gaelle.fromont-hankard@univ-tours.fr

Credit author statement

Jean-Philippe Antignac, Philippe Marchand, Ingrid Guiffard, Emmanuelle Bichon, Bruno le Bizec : OCP measurement

Sandy Figiel, Michèle Pinault, Karine Mahéo : Fatty acid analysis

Pascal Blanchet, Franck Bruyère, Romain Mathieu, Souhil Lebdaï, Georges Fournier, Jerome Rigaud, Luc Multigner : provide bioressources

Gaëlle Fromont, Luc Multigner : statistical analysis

Gaëlle Fromont, Jean-Philippe Antignac, Luc Multigner : manuscript writing

Gaëlle Fromont : study design

ABSTRACT

Although several studies have examined the relationship between organochlorine pesticides (OCPs) and prostate cancer (PCa) risk, no data are available concerning the association between OCPs concentrations in periprostatic adipose tissue (PPAT), which reflects cumulative exposure, and PCa aggressiveness. Moreover, no previous study has compared OCPs exposure in two distinct ethno-geographical populations. The objectives were to analyze OCPs in PPAT of PCa patients from either Mainland France or French West Indies in correlation with features of tumor aggressiveness, after adjusting for potential confounders such age, BMI, and polyunsaturated fatty acid (PUFA) content of PPAT.

PPAT was analyzed in 160 patients (110 Caucasians and 50 African-Caribbeans), 80 with an indolent tumor (ISUP group 1 + pT2), and 80 with an aggressive tumor (ISUP group more than 3 + pT3). The concentrations of 29 OCPs were measured in PPAT concomitantly with the characterization of PUFA content.

Exposure patterns of OCPs differed according to the ethno-geographical origin. Most OCPs were found at higher concentration in Caucasian patients, whereas pp'-DDE content was twice as high in African-Caribbeans. Chlordecone was only detected in PPAT from African-Caribbean patients. Most OCP concentrations were positively correlated with age, and some with BMI. After adjusting for age, BMI, and PUFA composition of PPAT, no significant association was found between OCPs content and risk of aggressive disease, except of mirex which appeared inversely associated with aggressive features of PCa in Caucasian patients.

These results highlight a significant ethno-geographic variation in internal exposure to OCPs, which likely reflects differences in consumption patterns. The inverse relationship observed between mirex concentration and markers of PCa aggressiveness need to be further investigated.

Keywords: prostate cancer; persistent organic pollutants; organochlorine pesticides; exposure; adipose tissue; fatty acids

1. INTRODUCTION

Prostate cancer (PCa) is the most frequently diagnosed cancer in men in Western countries. Significant ethnic disparities in PCa risk have been reported, with a higher incidence among African-American men compared with European-Americans (1). Similarly, in the French West Indies, where at least 90% of the population is of African origin, the incidence of PCa is twice as high as in the rest of France (2). In addition to genetic parameters, it has been suggested that environmental factors including dietary intake, play a role in the risk and progression of PCa (3). We have previously shown that the polyunsaturated fatty acid (PUFA) composition of peri-prostatic adipose tissue (PPAT), which reflects PUFA intake, was associated with PCa aggressiveness, and differed according to the ethno-geographical origin. Linoleic acid (LA), an essential n-6 PUFA, was 2-fold higher in African-Caribbean PPAT than in Caucasian patients, and was inversely associated with features of PCa aggressiveness. In Caucasians, a negative association was observed between eicosapentaenoic acid (EPA) levels (an n-3 PUFA) and disease aggressiveness (4).

Organochlorine pesticides (OCPs) are a group of Persistent Organic Pollutants (POPs) with endocrine disrupting and bioaccumulative properties (5). OCPs have been widely used in agriculture and pest control, and most of them have now been banned for decades in Western countries, due to public health concerns. However, due to their persistent nature, they are still present in the environment, food, and consequently in human matrices (6). Food and drinking water are the main sources of contamination (7). In the human body, due to their highly lipophilic properties, they tend to accumulate in tissues with high fat content, mainly adipose tissue (8).

Several epidemiologic studies have investigated the relationship between OCP exposure levels and PCa risk, both in occupational exposure and in the general population, with mainly tenuous, inconsistent or opposite findings (9, 10, 11, 12). Furthermore, some studies have focused on the association between exposure to selected OCPs and aggressive disease in the general population, reporting significant associations of high risk or metastatic PCa, with p-p'DDE, oxychlorane, and chlordecone, respectively (13-16). In most of these studies, exposure to OCPs was evaluated either by questionnaires or by blood measurements. Because OCPs accumulates in fat, their measurement in adipose tissue represent cumulative internal exposure, accounting for all exposure routes and sources (17). Therefore, adipose tissue, when available, is a good indicator of long-term exposure for lipophilic chemicals.

In the present study, we aimed to determine the concentration levels of 29 OCPs and related metabolites from the Stockholm convention list, in PPAT from patients with PCa. The measurements

93 were performed in two populations of different ethno-geographical origin, and in association with
94 features of disease aggressiveness, after adjusting or not for PUFA composition.

95

Journal Pre-proof

2. MATERIAL AND METHODS

2. 1. Studied population and collected biological tissues

PPAT has been sampled during the surgical procedure in patients treated by radical prostatectomy for clinically localized PCa in 6 University Hospitals. Five of these centers mainly recruited Caucasian patients, while the one located in the French West Indies recruited African-Caribbean patients. All patients have signed an informed consent, after agreement of the Ethical Committee/IRB, clinicaltrials registration number NCT03214315. Clinical and biological annotations included age, ethno-geographic origin, preoperative prostatic specific antigen (PSA), pathological stage, ISUP group, body mass index (BMI). From 1000 cases included in the database, patients with aggressive PCa have been selected on the criteria of high risk disease, i.e. ISUP group more than 3 and pT3 stage. Each of these patients have then been matched (within each center) for the age with one patient with indolent disease. Criteria for indolent PCa include ISUP group 1 and pT2 stage and preoperative PSA less than 10 ng/ml. At the end, PPAT from 160 patients (110 Caucasians, 50 African-Caribbeans) have been analyzed, including 80 with indolent disease and 80 with aggressive disease. The characteristics of this studied population are summarized in table 1.

2. 2. Organochlorine pollutants (OCP) extraction and quantification by gas chromatography-high resolution mass spectrometry (GC-HRMS) and liquid chromatography-tandem mass spectrometry (LC-MS/MS)

Sample preparation for adipose tissues consisted in 1 g freeze-dried sample extracted with a toluene/acetone mixture (70:30, v/v) using a Speed Extractor E-914 from Buchi (Rungis, France). Pressure and temperature were set at 100 bars and 120 °C respectively.

Tissue extracts were then purified on a Gel Permeation Chromatography (GPC) system. GPC extracts were split into two fractions. The first fraction was used for OCP analysis by GC-HRMS, the second for chlordecone analysis by LC-MS/MS. Extracts were carefully evaporated to dryness under a gentle stream of nitrogen before reconstituting the chlordecone fraction in 50 µL acetonitrile and the OCP fraction in 50 µL of ¹³C-PCB-111 at 20 pg.L⁻¹ (in nonane for GC-MS analysis).

The methodologies applied to isolate, detect, and quantify the targeted organochlorine pesticides except chlordecone, using GC-HRMS, have been described earlier (12). Briefly, ¹³C-labeled congeners were added to each sample for quantification according to the isotopic dilution method. OC measurements were performed by gas chromatography (Agilent 7890A) coupled to high-resolution mass spectrometry (GC-HRMS) on electromagnetic sector instruments (JEOL MS 700D or 800D), operating at 10,000 resolution and in the single ion monitoring (SIM) acquisition mode. QA/QC

procedures included systematic analysis of negative (procedural blank, n = 3) and positive (standard reference compounds, n = 1) quality control samples in each batch of analysis. All these methods were fully validated .

The methodology applied to isolate, detect, and quantify chlordecone using LC–MS/MS has been described earlier (18). An Acquity Ultra Performance Liquid Chromatography (UPLC) System (Waters, Milford, MA, USA) equipped with a C30 Thermo Accucore column (100 mm × 2.1 mm i.d., 2.6 µm granulometry) was used for liquid chromatography (LC) separation. The column temperature was set at 40 °C. A Xevo TQ-S triple quadrupole mass spectrometer (Waters, Milford, MA, USA) equipped with an electrospray ionization source was used in negative mode (ESI-). TargetLynx V 4.1 (Waters, Milford, MA, USA) software was used for the integration and assignment of all chromatographic peaks acquired in the selected reaction monitoring (SRM) mode.

The following OCPs were analyzed: hexachlorobenzene (HCB), pentachlorobenzene (PeCB), α -hexachlorocyclohexane (α -HCH), β -hexachlorocyclohexane (β -HCH), γ -hexachlorocyclohexane (γ -HCH), δ -hexachlorocyclohexane (δ -HCH), α -chlordane, γ -chlordane, cis-nonachlor, trans-nonachlor, heptachlor, cis-heptachlor-epoxyde, trans-heptachlor-epoxyde, oxychlordane, α -endosulfan, β -endosulfan, endosulfan-sulfate, aldrin, endrin, dieldrin, o,p'-DDT, p,p'-DDT, o,p'-DDD, p,p'-DDD, o,p'-DDE, p,p'-DDE, methoxychlor, mirex, chlordecone. Results are expressed as ng per g of extracted lipids.

2. 3. Fatty acid analysis of PPAT

Total lipids were extracted, according to the Folch method , with chloroform/methanol 2:1 and saline solution. Organic phase was collected and triglycerides were purified by preparative Thin Layer Chromatography. FA were transmethylated with 14% boron trifluoride in methanol, and FA methyl esters were dissolved in hexane and analyzed by capillary gas chromatography. FA were identified by comparison of their retention times with those of a standard mixture (Supelco™ 37 component FA methyl ester MIX, Supelco). Results for each FA expressed as mole % of total chromatogram fatty acids area.

2. 4. Statistical analysis

Statistical analyses were carried out with Addinsoft™ XLSTAT (New York, NY, USA). Comparison between groups was performed using the nonparametric Mann-Whitney U test. The relationship between 2 continuous variables was evaluated using the nonparametric Spearman rank correlation test. Logistic regressions were performed to evaluate the association between each OCP and features of aggressive versus indolent disease. OCPs were entered in the logistic models as 2 categories,

above and below the median value. Each model was adjusted on age and BMI as continuous variables, because OCPs are known to bio-accumulate. The second model also adjusted for EPA and AL content of PPAT, PUFA that have been associated with features of PCa aggressiveness, in the Caucasian and African-Caribbean population, respectively, and that are therefore confounders (4). The p values have not been corrected for multiplicity, because the study has been performed for descriptive purposes, focused on a few planned comparisons. Therefore the following results should clearly be labelled as exploratory.

3. RESULTS

3. 1. OCP concentration levels in PPAT

The following OCPs were found below the limit of detection (LOD) in more than 70% of individuals in both group of patients: α -HCH, γ -HCH, δ -HCH, α -chlordane, γ -chlordane, heptachlor, trans-heptachlor-epoxyde, α -endosulfan, β -endosulfan, endosulfan-sulfate, aldrin, endrin, o,p'-DDT, o,p'-DDD, o,p'-DDE, methoxychlor.

The following OCPs were detectable in all patients in both groups: HCB, PeCB, β -HCH, trans-nonachlor, cis-heptachlor-epoxyde, oxychlordane, dieldrin, p,p'-DDE, mirex.

Cis-nonachlor, p,p'-DDT and p,p'-DDD were detectable in more than 70% of patients in both groups. Chlordane was detectable in around 70% of patients in PPAT from African-Caribbean patients, and below the LOD in PPAT from Caucasian patients. For the few cases below the LOD, non-detected values have been replaced by limits of detection (upper bound approach).

The main contributors to the OCP contamination profile observed in PPAT are p, p'-DDE, β -HCH, trans-nonachlor, HCB, and oxychlordane (figure 1).

3. 2. The OCP exposure patterns in PPAT differed according to the ethno-geographical origin

Most OCPs were found at higher concentration level in PPAT from Caucasian patients when compared to PPAT from African-Caribbean patients, including β -HCH, trans-nonachlor, HCB, PeCB, oxychlordane, dieldrine, cis-heptachlor-epoxyde, and cis-nonachlor (table 2). In contrast, OCP concentrations in PPAT were higher in African-Caribbean patients for p,p'-DDE (more than twice), p,p'-DDT, and mirex. Chlordane was detected only in PPAT from African-Caribbean patients.

3. 3. Correlations between OCPs

Spearman's rank correlations between OCPs are represented in figure 2 as heatmaps built for each population. Strong correlations were observed between HCB, β -HCH, cis-nonachlor, trans-nonachlor, cis-heptachlor-epoxyde, oxychlordane, dieldrin, and p,p'-DDE, regardless of the ethno-geographic group. PeCB was slightly (but significantly) correlated with cis-nonachlor, trans-nonachlor, cis-heptachlor-epoxyde, oxychlordane, dieldrin, and mirex, but only in Caucasian patients.

In the Caucasian population, mirex appeared significantly correlated with HCB, PeCB, β -HCH, cis-nonachlor, trans-nonachlor, cis-heptachlor-epoxyde, oxychlordane, and dieldrin. In African-Caribbean patients, mirex was found significantly correlated with β -HCH, p,p'-DDE, and chlordane.

3. 4. Correlation between OCP accumulation in PPAT, age and BMI

In Caucasian patients, most OCP concentrations were positively correlated with patient's age, including p, p'-DDE (Rho: 0.264, p=0.006, Spearman test), β -HCH (Rho: 0.367, p=0.0001), trans-nonachlor (Rho: 0.409, p<0.0001), HCB (Rho: 0.303, p=0.0015), oxychlordane (Rho: 0.411, p<0.0001), mirex (Rho: 0.362, p=0.0002), dieldrin (Rho: 0.373, p<0.0001), cis-heptachlor-epoxyde (Rho: 0.400, p<0.0001), and PeCB (Rho: 0.221, p=0.02). In African-Caribbean patients, only p, p'-DDE and oxychlordane increased statistically with patient's age (Rho: 0.358 and 0.308, p=0.01 and 0.03, respectively).

In Caucasian patients, the following OCPs were positively correlated with the BMI: β -HCH (Rho: 0.251, p=0.009), dieldrin (Rho: 0.219, p=0.02), and cis-heptachlor-epoxyde (Rho: 0.225, p=0.02). Mirex concentration in PPAT from Caucasian patients was associated with lower BMI (Rho: -0.251, p=0.009). In African-Caribbean patients, both dieldrin and chlordecone concentrations in PPAT were positively correlated with BMI (Rho: 0.343 and 0.381, p=0.07 and 0.008, respectively).

3. 5. OCP exposure and tumor aggressiveness

The association between OCP internal exposure levels and features of aggressive disease compared to indolent disease has been analyzed using logistic regression models adjusted for age and BMI (tables 3 and 4). Moreover, since EPA and AL were associated with aggressive disease in Caucasians and African-Caribbeans, respectively (4), they have been included in a second model (tables 3 and 4). Results concerning omega 6 and omega 3 fatty acids content in PPAT are available in the supplemental table. In the Caucasian population, mirex was the only OCP significantly associated with features of aggressive disease, with an inverse association after adjustment for age and BMI (p=0.027), and after adjustment for age, BMI, and EPA content (p=0.029). In the African-Caribbean population, the inverse association between mirex and features of PCa aggressiveness was close to significance (p=0.06 and 0.10 in models 1 and 2, respectively). In this population, we also found almost significant associations between aggressive disease and oxychlordane, dieldrine and cis heptachlorepoxyde, but only in the first model.

4. DISCUSSION

Most of the considered OCPs have been withdrawn from the market for a long time, but their low ability to degrade leads to their persistence in the environment as well as their bioaccumulation. As a result, populations are still exposed, mainly through dietary intake. Measurement of OCPs and their metabolites in biological matrices is considered a reference method to assess exposure at steady state. Although most previous reports have analyzed OCPs content in blood, few studies have reported OCPs composition in adipose tissue, and none in PPAT. The concentration we found for p,p'-DDT and its metabolite p,p'-DDE is close or slightly higher to that described in populations of patients from Mainland France with either endometriosis (19) or gastric cancer (20), but is quite lower than those reported in Spanish or Danish women with breast cancer (21, 22), in non-cancer patients from Tunisia (23), and in PCa patients from Sweden (24). The differences observed could be due to the diversity of the populations studied, in terms of age, gender, geographical origin (and therefore exposure), and also in terms of location of the adipose tissue (subcutaneous versus deep fat). However, in all studies, the high ratio p,p'-DDE/p,p'-DDT reflects a preferential exposure to the metabolite and not to the parent molecule. Concerning the other OCPs, the concentration we found in the PPAT of Caucasian patients is of the same order as that reported in the most recent study performed in Mainland France (20).

We observed that most of OCPs concentrations increased with age, mainly in the Caucasian group, that reflects cumulative exposure, as previously reported (25). Some of them, including β -HCH, dieldrin, cis-heptachlor-epoxyde, and chlordecone in the African-Caribbean population, had concentrations increasing with the BMI. Although in some studies no significant relationship was found (24), high BMI has often been associated with higher POPs levels in adipose tissue (26). This association may be related to food intake, since individuals with a higher BMI are likely to have consumed more food and therefore more POPs. Alternatively, POPs are considered to have obesogenic effects through endocrine disruption (27). We observed herein that mirex is the only OCP which concentration in PPAT was inversely associated with BMI. Such an inverse relationship between BMI and mirex concentration has previously been reported (28). A potential explanation may be the dilution capabilities in adipose tissue of mirex, one of the most lipophilic compounds.

In the present study, we found a differential OCPs concentration in PPAT according to the ethno-geographical origin of the patients. β -HCH, trans-nonachlor, HCB, PeCB, oxychlordane, dieldrine, cis-heptachlor-epoxyde, and cis-nonachlor levels were higher in PPAT from Caucasian patients, whereas the concentrations of p,p'-DDE, p,p'-DDT, and mirex were higher in African-Caribbean patients. No previous study has compared the impregnation levels by OCPs in 2 distinct ethno-geographic

populations. However, the report of the serum concentration of OCPs in the Guadeloupean population by the Kannari Study (29) evidenced lower β -HCH and higher p,p'-DDE when compared to the levels identified in Mainland France by the French National Nutrition and Health Study (ENNS) (30). Since food is the main vector of exposure to OCPs, differences between Guadeloupe and Mainland France probably reflect differences in eating behaviors, but also differences in the geographical origins of food resources. In the West Indies, although some of the foodstuffs come from Mainland France, many others come directly from the USA, Central America, and other islands of the Caribbean. We found a pp'-DDE concentration twice as high in PPAT of African-Caribbean patients compared to Caucasian patients. Such differences in pp'-DDE blood levels have already been reported between different parts of the globe (31). The difference in p,p'-DDE between African-Caribbean and Caucasian men could be partly explained by more recent exposure in West Indies, due to the sporadic use of DDT for disease vector control. Pesticides HCB, β -HCH, cis-nonachlor, trans-nonachlor, cis-heptachlor-epoxyde, oxychlordan, and dieldrin, that we found well correlated to each other and that are in higher concentrations in PPAT from Caucasian patients, could come mainly from foodstuffs of Mainland France origin. Mirex was discreetly used in Guadeloupe to control the fire ant, and has also been worldwide used as a flame retardant. The greatest concentration in African-Caribbean PPAT could be explained by the geographical origins of food resources and perhaps to a lesser extent by the use of the molecule in the West Indies until early 1980s. Chlordane is an organochlorine insecticide used in the French West Indies between 1972 and 1993 to control banana weevil. This highly stable, persistent organic substance detected in soils is capable of contaminating plant or animal foodstuffs as well as aquatic environments (32). As expected, we detected chlordane only in PPAT from African-Caribbean patients. Furthermore, chlordane is poorly correlated with other contaminants, that clearly reflects its local food origin. We note that the only molecules for which a certain correlation is observed, are β -HCH, dieldrin and mirex, pesticides previously used in the French West Indies.

Several previous non occupational case-control studies have analyzed in the general population the relationship between PCa incidence and exposure to OCPs, determined by blood measurements. Although a meta-analysis failed to evidence an association between specific OCPs and increased incidence of PCa (10), individual studies led to conflicting results. Some reports found no significant associations (33-35), while others observed a positive association between PCa incidence and several OCPs, including β -HCH (36, 37), oxychlordan (38), and p,p'-DDE (13). In another study, oxychlordan has also been found to be correlated with the risk of metastatic PCa (14). One case-control study has to date analyzed the link between chlordane plasma concentrations and PCa in a Guadeloupean population, and the authors described a significant association with a dose-response positive

relationship (15). The follow-up of PCa patients included in this case-control study also evidenced a link between risk of biochemical recurrence after surgical treatment and plasma chlordecone concentration (16). Except for these above mentioned studies, most of the previous reports have focused on the association between OCPs and total PCa risk. In the present study, we used indolent cases as the reference group, that allowed to identify factors associated with aggressiveness, defined by both ISUP group more than 3 and pT3 stage. In the African-Caribbean group only, we found associations close to significance between features of aggressive disease and oxychlorodane, dieldrine and cis heptachlorepoxyde. However, this almost significance was lost when AL content of PPAT was entered into the model, PUFA that we previously found to be inversely associated with PCa aggressiveness, and which is therefore a confounder (4). We also observed an inverse association between mirex content in PPAT and aggressive PCa. This association was significant in the Caucasian group only, and remained significant after adjusting for age, BMI, and for EPA content in PPAT. (4). In the African-Caribbean population, the inverse association between mirex and features of aggressiveness was close to significance, after adjusting or not for PUFA content. This inverse association has already been observed with other types of pesticides, such as polychlorinated biphenyls (PCBs) (13, 14). Moreover, the case-control study performed in a Canadian population also evidenced an almost significant inverse association between plasma mirex levels and PCa risk (34). Although it is unexpected that mirex could be protective for PCa, it is possible that this inverse association could reflect hormonal modifications linked to mirex exposure, and this finding warrants further studies. An inverse association between testosterone levels and mirex serum concentrations has previously been described in boys and men, after adjusting for potential confounders such age and BMI (39, 40). Also, we cannot exclude that this inverse association could result from bias due to either unmeasured confounders, or socio-demographic variables that could impact both dietary intake or PSA screening.

The limit of the present study is first the low number of cases, particularly in the African-Caribbean population, that limits its power. Moreover, except for the ethnic origin, genetic determinants of PCa aggressiveness, including family history of PCa, were not available and therefore were not taken into account.

The study has nevertheless several strengths: first, OCPs concentrations were measured directly in adipose tissue, the preferential reservoir of lipophilic pesticides accumulation in the human body, and the most appropriate tissue for assessing long term exposure. In addition, the use of indolent PCa cases as a reference group exempts the study from the bias linked to the identification of control cases. Indeed, some OCPs, and particularly p-p'DDE, the main contributor of OCPs profile in PPAT, are known inhibitors of the androgen receptor, and have been shown to repress PSA levels in human

PCa cell lines at high concentration (41). Strong exposure to these OCPs may therefore produce a false-negative PSA test when screening for prostate cancer.

In conclusion, the present study showed a differential OCPs internal exposure levels and patterns in adipose tissue according to the ethno-geographical origin. The inverse relationship in adjusted models between mirex concentration and features of PCa aggressiveness deserves further mechanistic and epidemiological studies.

REFERENCES

- 1- Evans S, Metcalfe C, Ibrahim F, Persad R, Ben-Shlomo Y. Investigating Black-White differences in prostate cancer prognosis: A systematic review and meta-analysis. *Int J Cancer* 2008;123(2):430-5.
- 2- Brureau L, Multigner L, Wallois A, Verhoest G, Ndong JR, Fofana M, Blanchet P. [Prostate cancer in Guadeloupe (French West Indies): incidence, mortality and clinicopathological features]. *Bull Cancer* 2009; 96(2): 165-70.
- 3- Sonn GA, Aronson W, Litwin MS. Impact of diet on prostate cancer: a review. *Prostate Cancer Prostatic Dis* 2005; 8(4):304-10.
- 4- Figiel S, Pinault M, Domingo I, Guimaraes C, Guibon R, Besson P, Tavernier E, Blanchet P, Multigner L, Bruyère F, Haillot O, Mathieu R, Vincendeau S, Rioux-Leclercq N, Lebdaï S, Azzouzi AR, Perrouin-Verbe MA, Fournier G, Doucet L, Rigaud J, Renaudin K, Mahéo K, Fromont G. Fatty acid profile in peri-prostatic adipose tissue and prostate cancer aggressiveness in African-Caribbean and Caucasian patients. *Eur J Cancer* 2018; 91: 107-15.
- 5- McKinley R, Plant JA, Bell JN, Voulvoulis N. Endocrine disrupting pesticides: implications for risk assessment. *Environ Int* 2008; 34: 168-83.
- 6- Patterson DG, Wong LY, Turner WE, Caudill SP, Dipietro ES, McClure PC, Cash TP, Osterloh JD, Pirkle JL, Samson EJ, Needham LL. Levels in the US population of those persistent organic pollutants (2003-2004) included in the Stockholm convention or in other long range transboundary air pollution agreements. *Environ Sci Technol* 2009; 43: 1211-8.
- 7- Schecter A, Colacino J, Haffner D, Patel K, Opel M, Papke O, et al. Perfluorinated compounds, polychlorinated biphenyls, and organochlorine pesticides contamination in composite food samples from dallas, Texas, USA. *Environ Health Perspect* 2010; 118: 796-802.
- 8- Yilmaz B, Terekci H, Sandal S, Kekestimur F. Endocrine disrupting chemicals: exposure, effects on human health, mechanism of action, models for testing and strategies for prevention. *Rev Endoc Metab disord* 2020; 21: 127-47.
- 9- Lewis-Mikhael AM, Bueno-Cavanillas A, Ofir Guiron T, Olmedo-Requena R, Delgado-Rodriguez M, Jimenez-Moleon JJ. Occupational exposure to pesticides and prostate cancer: a systematic review and meta-analysis. *Occup Environ Med* 2016; 73: 134-44.

- 10- Lewis-Mikhael AM, Olmedo-Requena R, Martinez-Ruiz V, Bueno-Cavanillas A, Jimenez-Moleon JJ. Organochlorine pesticides and prostate cancer, is there an association? A meta-analysis of epidemiological evidence. *Cancer Causes Control* 2015; 26: 1375-92.
- 11- Koutros S, Beane Freeman LE, Lubin JH, Heltshe SL, Andreotti G, Barry KH, DellaValle CT, Hoppin JA, Sandler DP, Lynch CF, Blair A, Alavanja MC. Risk of total and aggressive prostate cancer and pesticide use in the Agricultural Health Study. *Am J Epidemiol.* 2013; 177(1): 59-74.
- 12- Pardo LA, Beane Freeman LE, Lerro CC, Andreotti G, Hofmann JN, Parks CG, Sandler DP, Lubin JH, Blair A, Koutros S. Pesticide exposure and risk of aggressive prostate cancer among private pesticide applicators. *Environ Health.* 2020; 19(1): 30.
- 13- Emeville E, Giusti A, Coumoul X, et al. Associations of plasma concentrations of dichlorodiphenyldichloroethylene and polychlorinated biphenyls with prostate cancer: a case-control study in Guadeloupe (French West Indies). *Environ Health Perspect* 2015; 123 : 317-23.
- 14- Koutros S, Langseth H, Grimsrud TK, Barr DB, Vermeulen R, Portengen L, Wacholder S, Freeman LE, Blair A, Hayes RB, Rothman N, Engel LS. Prediagnostic Serum Organochlorine Concentrations and Metastatic Prostate Cancer: A Nested Case-Control Study in the Norwegian Janus Serum Bank Cohort. *Environ Health Perspect.* 2015; 123(9): 867-72.
- 15- Multigner L, Ndong JR, Giusti A, Romana M, Delacroix-Maillard H, Cordier S, Jégou B, Thome JP, Blanchet P. Chlordecone exposure and risk of prostate cancer. *J Clin Oncol.* 2010; 28(21): 3457-62.
- 16- Brureau L, Emeville E, Helissey C, Thome JP, Multigner L, Blanchet P. Endocrine disrupting-chemicals and biochemical recurrence of prostate cancer after prostatectomy: A cohort study in Guadeloupe (French West Indies). *Int J Cancer.* 2020 Feb 1;146(3):657-663.
- 17- Kohlmeier L., Kohlmeier M. Adipose tissue as a medium for epidemiologic exposure assessment. *Environ. Health Perspect* 1995; 103, S3: 99-106.
- 18- Bichon E, Guiffard I, Venisseau A, Marchand P, Antignac JP, Le Bizec B. Ultratrace quantification for chlordecone in human fluids and tissues. *J Chromatography A* 2015; 1408: 169-77.

- 19- Ploteau S, Antignac JP, Volteau C, Marchand P, Venisseau A, Vacher V, Le Bizec B. Distribution of persistent organic pollutants in serum, omental, and parietal adipose tissue of french women with deep infiltrating endometriosis and circulating versus stored ratio as new marker of exposure. *Environ Int* 2016; 97: 125-36.
- 20- Perrot-Appianat M, Pimple C, Cano-sanchez G, Antignac JP, Pocard M. Detection of persistent organic pollutants in omental adipose tissue from patients with diffuse gastric cancer: a pilot study. *Cancers* 2021; 13: 4874.
- 21- Artacho-Cordon F, Fernandez-Rodriguez M, Garde C, Salamanca E, Iribarne-Duran LM, Torné P, Exposito J, Papay-Ramirez L, Fernandez MF, Olea N, Arrebola JP. Serum and adipose tissues as matrices for assessment of exposure to persistent organic pollutants in breast cancer patients. *Environ Res* 2015; 142: 633-43.
- 22- Roswall N, Sorensen M, Tjønneland A, Raaschou-Nielsen O. Organochlorine concentrations in adipose tissue and survival in postmenopausal danish breast cancer patients. *Environ Res* 2018; 163: 237-48.
- 23- Achour A, Derouiche A, Barhoumi B, Kort B, Cherif D, Bouabdallah S, Sakly M, Ben Rhouma K, Touil S, Ridha Driss M, Tebourbi O. Organochlorine pesticides and polychlorinated biphenyls in human adipose tissue from northern Tunisia: current extent of contamination and contributions of sociodemographic characteristics and dietary habits. *Environ Res* 2017; 156: 635-43.
- 24- Hardell L, Andersson SO, Carlberg M, Bohr L, van Bavel B, Lindstrom G, Bjornfoth H, Ginman C. Adipose tissue concentrations of persistent organic pollutants and the risk of prostate cancer. *JOEM* 2006; 48: 700-7.
- 25- Li M, Wang R, Su C, Li J, Wu Z. Temporal Trends of Exposure to Organochlorine Pesticides in the United States: A Population Study from 2005 to 2016. *Int J Environ Res Public Health*. 2022; 19(7): 3862.
- 26- Sousa S, Maia ML, Delerue-Matos C, Calhau C, Dominguez VF. The role of adipose tissue analysis on environmental pollutants biomonitoring in women : the european scenario. *Sci Total Environ*. 2022; 806: 150922.

- 27- Aaseth J, Javorac D, Djordjevic AB, Bulat Z, Skalny AV, Zaitseva IP, Aschner M, Tinkov AA. The Role of Persistent Organic Pollutants in Obesity: A Review of Laboratory and Epidemiological Studies. *Toxics*. 2022; 10(2): 65.
- 28- Kanazawa A, Miyasita C, Okada E, Kobayashi S, Washino N, Sasaki S, Yoshioka E, Mizutani F, Chisaki Y, Saijo Y, Kishi R. Blood persistent organochlorine pesticides in pregnant women in relation to physical and environmental variables in The Hokkaido Study on Environment and Children's Health. *Sci Total Environ*. 2012; 426: 73-82.
- 29- Dereumeaux C, Saoudi A, Guldner L, Pecheux M, Chesneau J, Thomé JP, Ledrans M, Tertre AL, Denys S, Fillol C. Chlordecone and organochlorine compound levels in the French West Indies population in 2013-2014. *Environ Sci Pollut Res Int*. 2020; 27(33): 41033-41045.
- 30- Saoudi A, Fréry N, Zeghnoun A, Bidondo ML, Deschamps V, Göen T, Garnier R, Guldner L. Serum levels of organochlorine pesticides in the French adult population: the French National Nutrition and Health Study (ENNS), 2006-2007. *Sci Total Environ*. 2014; 472: 1089-99.
- 31- Koureas M, Rousou X, Haftiki H, Mouchtouri VA, Rachiotis G, Rakitski V, Tsakalov A, Hadjichristodoulou C. Spatial and temporal distribution of pp'-DDE blood levels across the globe. A systematic review and meta-analysis. *Science of the total environment* 2019; 686: 440-451.
- 32- Cabidoche YM, Achard R, Cattani P, Clermont-Dauphin C, Massat F, Sansoulet J. Long-term pollution by chlordecone of tropical volcanic soils in the French West Indies: a simple leaching model accounts for current residue. *Environ Pollut*. 2009; 157(5): 1697-705.
- 33- Sawada N, Iwasaki M, Inoue M, Itoh H, Sasazuki S, Yamaji T, Shimazu T, Tsugane S; Japan Public Health Center Based Prospective (JPHC) Study Group. Plasma organochlorines and subsequent risk of prostate cancer in Japanese men: a nested case-control study. *Environ Health Perspect*. 2010; 118(5): 659-65.
- 34- Aronson KJ, Wilson JW, Hamel M, Diarsvitri W, Fan W, Woolcott C, Heaton JP, Nickel JC, Macneily A, Morales A.J. Plasma organochlorine levels and prostate cancer risk. *Expo Sci Environ Epidemiol*. 2010; 20(5): 434-45.
- 35- Lim JE, Nam C, Yang J, Rha KH, Lim KM, Jee SH. Serum persistent organic pollutants (POPs) and prostate cancer risk: A case-cohort study. *Int J Hyg Environ Health*. 2017; 220(5): 849-856.

- 36- Kumar V, Yadav CS, Singh S, Goel S, Ahmed RS, Gupta S, Grover RK, Banerjee BD. CYP 1A1 polymorphism and organochlorine pesticides levels in the etiology of prostate cancer. *Chemosphere*. 2010; 81(4): 464-8.
- 37- Xu X, Dailey AB, Talbott EO, Ilacqua VA, Kearney G, Asal NR. Associations of serum concentrations of organochlorine pesticides with breast cancer and prostate cancer in U.S. adults. *Environ Health Perspect*. 2010; 118(1): 60-6.
- 38- Ritchie JM, Vial SL, Fuortes LJ, Guo H, Reedy VE, Smith EM. Organochlorine and risk of prostate cancer. *JOEM* 2003; 45: 692-702.
- 39- Araki A, Miyashita C, Mitsui T, Goudarzi H, Mizutani F, Chisaki Y, Itoh S, Sasaki S, Cho K, Moriya K, Shinohara N, Nonomura K, Kishi R. Prenatal organochlorine pesticide exposure and the disruption of steroids and reproductive hormones in cord blood: the Hokkaido study. *Environment International* 2018; 110: 1-13.
- 40- Freire C, Koifman RJ, Sarcinelli PN, Rosa ACS, Clapauch R, Koifman S. Association between serum levels of organochlorine pesticides and sex hormones in adults living in a heavily contaminated area in brazil. *Int J Hyg Env Health* 2014; 217: 370-378.
- 41- Wong LI, Labrecque MP, Ibuki N, Cox ME, Elliott JE, Beischlag TV. p,p'-Dichlorodiphenyltrichloroethane (p,p'-DDT) and p,p'-dichlorodiphenyldichloroethylene (p,p'-DDE) repress prostate specific antigen levels in human prostate cancer cell lines. *Chem Biol Interact*. 2015; 230:40-9.

FIGURE LEGENDS

Figure 1: Main contributors to the exposure profiles in PPAT in France mainland and in Guadeloupe.

Figure 2: Heatmaps displaying the Spearman's rank correlation coefficients between OCPs in the African-Caribbean (A) and Caucasian (B) populations. The color relates to the direction of the association (blue: positive, red: negative), and the intensity to the correlation coefficient magnitude.

489 **ACKNOWLEDGMENT**

490 The authors thank "Canceropole Grand Ouest", "Region Bretagne", "Region Pays-de-Loire" and
491 "Region "Centre-Val-de-Loire", for financial support


492

493


494

495


Journal Pre-proof




A




B



A



B



Tables

Table 1. Characteristics of the studied population.

	Indolent PCa		<i>p</i>	Aggressive PCa		<i>p</i>
	Caucasians (n = 55)	AC (n = 25)		Caucasians (n = 55)	AC (n = 25)	
Pre-operative PSA, median (range)	6.4 ng/ml (1.1–9.9)	5.7 ng/ml (0.3–9)	0.07	9.4 ng/ml (3.5–30)	9.2 ng/ml (3.7–30)	0.87
Age, median (range)	64 y (50–77)	64 y (44–75)	0.62	66 y (46–75)	65 y (48–76)	0.71
BMI, median (range)	26 (18.5–39.1)	26.8 (21.9–38.4)	0.34	26.1 (20.3–34)	25.6 (20.6–32.9)	0.63

AC: African-Caribbeans; BMI: Body mass Index.

Table 2. OCP internal concentration levels (ng/g lipid weight) measured in PPAT from Caucasian and African-Caribbean, in both indolent and aggressive groups.

OCP	LOD (ng/g lw)	Ethno-geographic group	n (%) <LOD	Concentration (ng/g lw) Median (range)	p
p,p'-DDE	0.32	Caucasian	0/110 (0%)	124.6 (16.6–1228.8)	<0.0001
		African-Caribbean	0/50 (0%)	277 (45.7–3828.4)	
β HCH	0.42	Caucasian	0/110 (0%)	38.5 (4.3–229.8)	0.04
		African-Caribbean	0/50 (0%)	26.1 (2.6–375.0)	
Trans-nonachlor	0.28	Caucasian	0/110 (0%)	15.7 (3.9–77.5)	<0.0001
		African-Caribbean	0/50 (0%)	9.7 (1.5–35.4)	
HCB	0.05	Caucasian	0/110 (0%)	14.7 (4.6–41.3)	<0.0001
		African-Caribbean	0/50 (0%)	7.2 (2.0–21.5)	
Mirex	0.06	Caucasian	0/110 (0%)	3.2 (0.86–14.5)	<0.0001
		African-Caribbean	0/50 (0%)	7.4 (2.8–150.9)	
Oxychlordane	0.22	Caucasian	0/110 (0%)	14.0 (3.7–38.8)	<0.0001
		African-Caribbean	0/50 (0%)	6.0 (1.0–28.6)	
Dieldrine	0.17	Caucasian	0/110 (0%)	10.5 (2.8–45.7)	<0.0001
		African-Caribbean	0/50 (0%)	3.7 (0.57–45.4)	
Cis-heptachlor-epoxyde	0.08	Caucasian	0/110 (0%)	7.9 (2.7–43.4)	<0.0001
		African-Caribbean	0/50 (0%)	3.0 (0.58–36.3)	
p, p'-DDT	0.54	Caucasian	23/110 (20.9%)	2.7 (0.54–17.3)	0.004
		African-Caribbean	14/50 (28%)	4.2 (0.70–105.0)	
Cis-nonachlor	0.28	Caucasian	0/110 (0%)	2.8 (0.45–14.8)	<0.0001
		African-Caribbean	9/50 (18%)	1.5 (0.28–4.2)	
p, p'-DDD	0.14	Caucasian	17/110 (15.4%)	0.66 (0.14–4.4)	0.09
		African-Caribbean	3/50 (6%)	0.73 (0.23–17.1)	
PeCB	0.05	Caucasian	0/110 (0%)	0.29 (0.05–2.2)	<0.0001
		African-Caribbean	0/50 (0%)	0.17 (0.07–1.2)	
Chlordecone	0.08	Caucasian	110/110 (100%)	ND	<0.0001
		African-Caribbean	5/50 (10%)	0.23 (0.08–9.49)	

Table 3. Association between OCP internal concentration levels in PPAT from Caucasians patients, and risk of aggressive versus indolent disease. Logistic regression after adjustment for age and BMI (model 1) and after adjustment for age, BMI, and EPA (model 2).

OCP	OR Model 1 (95% CI)	P	OR Model 2 (95% CI)	P
p,p'-DDE	1.00 (ref)	0.94	1.00 (ref)	0.99
Below median	1.032 (0.472–2.254)		0.997 (0.435–2.285)	
Above median				
β HCH	1.00 (ref)	0.72	1.00 (ref)	0.33
Below median	0.868 (0.394–1.915)		0.648 (0.272–1.544)	
Above median				
Trans-nonachlor	1.00 (ref)	0.41	1.00 (ref)	0.38
Below median	0.713 (0.317–1.604)		0.670 (0.277–1.624)	
Above median				
HCB	1.00 (ref)	0.21	1.00 (ref)	0.09
Below median	1.645 (0.760–3.562)		2.129 (0.894–5.070)	
Above median				
Mirex	1.00 (ref)	0.027	1.00 (ref)	0.029
Below median	0.381 (0.161–0.898)		0.358 (0.142–0.900)	
Above median				
Oxychlorane	1.00 (ref)	0.69	1.00 (ref)	0.84
Below median	1.175 (0.535–2.580)		1.090 (0.468–2.539)	
Above median				
Dieldrine	1.00 (ref)	0.76	1.00 (ref)	0.84
Below median	0.881 (0.388–2.001)		0.915 (0.379–2.208)	
Above median				
Cis heptachlorepoxyde	1.00 (ref)	0.37	1.00 (ref)	0.71
Below median	1.454 (0.644–3.286)		1.185 (0.492–2.855)	
Above median				
p, p'-DDT	1.00 (ref)	0.58	1.00 (ref)	0.66
Below median	0.809 (0.380–1.720)		0.833 (0.370–1.876)	
Above median				
Cis-nonachlor	1.00 (ref)	0.12	1.00 (ref)	0.29
Below median	0.538 (0.243–1.188)	0.09	0.629 (0.266–1.492)	0.12
Above median	1.00 (ref)		1.00 (ref)	
PeCB	0.517 (0.242–1.102)		0.530 (0.236–1.192)	
Below median				
Above median				
p, p'-DDD	1.00 (ref)	0.81	1.00 (ref)	0.88
Below median	0.913 (0.431–1.936)		0.942 (0.419–2.120)	
Above median				

Table 4. Association between OCP internal concentration levels in PPAT from African-Caribbean patients, and risk of aggressive versus indolent disease. Logistic regression after adjustment for age and BMI (model 1) and after adjustment for age, BMI, and AL (model 2).

OCP	OR Model 1 (95%CI)	P	OR Model 2 (95% CI)	P
p,p'-DDE	1.00 (ref)	0.94	1.00 (ref)	0.92
<i>Below median</i>	1.048 (0.279–3.933)		1.089 (0.185–6.424)	
<i>Above median</i>				
β HCH	1.00 (ref)	0.11	1.00 (ref)	0.29
<i>Below median</i>	2.891 (0.789–10.593)		2.481 (0.466–13.217)	
<i>Above median</i>				
Trans-nonachlor	1.00 (ref)	0.14	1.00 (ref)	0.30
<i>Below median</i>	2.561 (0.745–8.806)		2.232 (0.486–10.255)	
<i>Above median</i>				
HCB	1.00 (ref)	0.48	1.00 (ref)	0.73
<i>Below median</i>	1.514 (0.469–4.891)		1.296 (0.302–5.563)	
<i>Above median</i>				
Mirex	1.00 (ref)	0.08	1.00 (ref)	0.10
<i>Below median</i>	0.298 (0.085–1.050)		0.264 (0.052–1.347)	
<i>Above median</i>				
Oxychlordanes	1.00 (ref)	0.08	1.00 (ref)	1.18
<i>Below median</i>	3.525 (0.982–12.657)		2.954 (0.610–14.300)	
<i>Above median</i>				
Dieldrine	1.00 (ref)	0.05	1.00 (ref)	0.64
<i>Below median</i>	3.794 (1.008–14.284)		1.462 (0.290–7.369)	
<i>Above median</i>				
Cis heptachlorepoxide	1.00 (ref)	0.05	1.00 (ref)	0.30
<i>Below median</i>	3.375 (1.075–13.255)		2.216 (0.485–10.119)	
<i>Above median</i>				
p, p'-DDT	1.00 (ref)	0.88	1.00 (ref)	0.39
<i>Below median</i>	1.112 (0.343–3.610)		1.995 (0.415–9.595)	
<i>Above median</i>				
Cis-nonachlor	1.00 (ref)	0.81	1.00 (ref)	0.89
<i>Below median</i>	1.154 (0.361–3.687)	0.27	1.108 (0.255–4.803)	0.52
<i>Above median</i>	1.00 (ref)		1.00 (ref)	
PeCB	1.933 (0.598–6.249)		01.611 (0.376–6.902)	
<i>Below median</i>				
<i>Above median</i>				
p, p'-DDD	1.00 (ref)	0.23	1.00 (ref)	0.45
<i>Below median</i>	0.481 (0.146–1.591)		0.552 (0.119–2.560)	
<i>Above median</i>				
Chlordane	1.00 (ref)	0.18	1.00 (ref)	0.86
<i>Below median</i>	0.446 (0.134–1.484)		0.875 (0.180–4.244)	
<i>Above median</i>				

HIGHLIGHTS

- Organochlorine pesticides (OCPs) concentration in periprostatic adipose tissue from men with prostate cancer differed according to the ethno-geographical origin, which likely reflects differences in consumption patterns.
- Most OCPs were found at higher concentration in PPAT from Caucasian patients. Chlordecone was only found in PPAT of African-Caribbean patients, with moreover a DDE concentration twice as high as in Caucasian patients.
- After adjusting for age, BMI, and PUFA content, mirex concentration in PPAT from Caucasian patients was inversely associated with features of aggressive disease.

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: