



HAL
open science

Larval pikeperch *Sander lucioperca* cannibals are more efficient predators on zebrafish *Danio rerio* than non-cannibals

Tatiana Colchen, Yannick Ledoré, Pascal Fontaine, Fabrice Teletchea, Alain Pasquet

► **To cite this version:**

Tatiana Colchen, Yannick Ledoré, Pascal Fontaine, Fabrice Teletchea, Alain Pasquet. Larval pikeperch *Sander lucioperca* cannibals are more efficient predators on zebrafish *Danio rerio* than non-cannibals. *Aquaculture*, 2023, 575, pp.739756. 10.1016/j.aquaculture.2023.739756 . hal-04197437

HAL Id: hal-04197437

<https://univ-angers.hal.science/hal-04197437v1>

Submitted on 6 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1 Larval pikeperch *Sander lucioperca* cannibals are more efficient predators on zebrafish *Danio*
2 *rerio* than non-cannibals.

3 T. Colchen ^{(a)*}, Y. Ledoré ^(a), P. Fontaine ^(a), F. Teletchea ^(a), A. Pasquet ^(a)

4

5 ^(a): University of Lorraine, INRAE, URAFPA, F-54000, Nancy, France

6

7 Corresponding author: tatiana.colchen@gmail.com

8 Phone : (+33) 2 41 73 52 43

9

10 E-mail addresses of co-authors:

11 yannick.ledore@univ-lorraine.fr

12 p.fontaine@univ-lorraine.fr

13 fabrice.teletchea@univ-lorraine.fr

14 alain.pasquet@univ-lorraine.fr

15

16 *present address: University of Angers, Biodiv'AG, F-49000, Angers, France

17 **Highlights**

18

- 19 • Cannibalism is an intra-specific predation resulting in massive mortality in fish
- 20 • Pikeperch is a predatory freshwater fish with precocious onset of piscivory
- 21 • Predatory behavioral sequence is more efficient in cannibals than in non-cannibals
- 22 • The shift to piscivory involved a change from a S-shape to an attack without S-shape
- 23 • Cannibalism is linked to the onset of an efficient piscivory behaviour

24

25 **Abstract**

26 Cannibalism is an intra-specific predation often resulting in massive mortality in intensive
27 cultured fish species. In piscivorous species, the onset of predation is inevitable. This study
28 explored the link between predation abilities and the onset of cannibalism in pikeperch
29 larvae, a top predator species, with a potential for the European inland aquaculture.
30 Cannibals and non-cannibals, aged between 17 to 34 days-post-hatching, were tested to
31 compare their predatory behavior using zebrafish and pikeperch larvae as prey. Each tested
32 fish was confronted with three other fish (pikeperch or zebrafish) ($n = 204$ tests) in an arena
33 ($20 \times 7 \times 4$ cm with 2 cm of water height, 280 mL) and their behavior was recorded for 20
34 minutes. The normal predatory behavioral sequence (orientation, approach, attack and
35 capture) was much more efficient in cannibals (calculated as the ratio of number of captures
36 on the number of attacks) than in non-cannibals in both the predation (0.31 ± 0.28 vs. $0.05 \pm$
37 0.15 ; $p_{\text{adjust}} < 0.0001$) and the cannibalism (0.21 ± 0.29 vs. 0.02 ± 0.08 ; $p_{\text{adjust}} = 0.0006$) tests.
38 The shift to a piscivorous diet involved a modification of the attack tactic, replacing a “S-
39 shape” by an attack without “S-shape” but with a tail propulsion. This second tactic
40 developed earlier in cannibals whereas non-cannibals are only able to perform the “S-shape”
41 attack used during the first days of life. Cannibalism was linked to the onset of an efficient
42 piscivory behavior.

43

44 **Keywords:** piscivory, intra-specific predation, behavior, freshwater carnivorous fish.

45 **1. Introduction**

46 Pikeperch *Sander lucioperca* has high potential for inland aquaculture diversification in
47 Europe (Lappalainen et al., 2003; Wang et al., 2009; Policar et al., 2019). The reproduction
48 control (Kucharczyk et al., 2007) and the bio-economic feasibility of its intensive rearing are
49 established (Steenfeldt and Lund, 2008; Steenfeldt et al., 2015). Its demand has been
50 strengthened by the strong decline of wild catches from approximately 50.000 tons in 1950
51 to 20.000 tons in 2020 (FAO, 2021). Currently, in Europe, the number of pikeperch farms has
52 increased and the production is estimated to 1000-1500 tons per year. Intracohort
53 cannibalism is one of the major bottleneck for the inland aquaculture of piscivorous percid
54 species particularly during the larval stages of pikeperch (Steenfeldt, 2015), walleye *Sander*
55 *vitreus* (Cuff, 1980) and perch *Perca fluviatilis* (Kestemont et al., 2003). For pikeperch,
56 cannibalistic predation can start a few days after hatching (between 14 and 17 dph) and
57 between individuals of similar size (9.3 ± 1.0 mm; Colchen et al., 2019). In pikeperch larvae,
58 cannibalism could induce up to 53 % at 37 days post-hatching (dph) of mortality in farm
59 (density: 17 larvae.L⁻¹, temperature: 18°C, feeding regime: 3 *Artemia* nauplii.mL⁻¹; Steenfeldt
60 et al., 2011). Cannibalism started as early as 14 dph (9.3 ± 1.0 mm total length (TL)) and was
61 always observed after 80 dph (Colchen et al., 2019).

62 Pikeperch is a predatory freshwater fish, which become piscivorous during its
63 development. Predation includes all behavioral items required to find, catch, kill and ingest
64 prey (Curio, 2012). Over time, a predator improves its searching tactics, visual and olfactory
65 prey recognition and attack strategies; conversely prey improve camouflage and avoidance
66 abilities. From an energetical perspective, predation has often been associated with the
67 optimal foraging theory (MacArthur and Pianka, 1966; Krebs and Davies, 2009). Considering
68 that cannibalism is an intra-specific predation (Polis, 1981), the transition between predator

69 and cannibal status could be viewed as a shift between a more generalist to a specialist
70 predator, and it should depend on biotic conditions as prey presence or density. When prey
71 are lacking or scarce, predator would target conspecifics as prey substitute. Among
72 pikeperch, predatory behavior increases progressively during ontogeny (Colchen et al.
73 2020a). The transition from embryonic stage to larval stage is characterized by the start of
74 exogeneous feeding (5.87 ± 0.77 mm TL; Ott et al., 2012), during which pikeperch feed first
75 on zooplankton with a typical predatory behavior called “S-shape” attack marked by a stop
76 of the fish and the formation of a “S” with his body before the propulsion towards prey
77 (Colchen et al., 2019; Houde, 2001). After that, pikeperch larvae progressively become
78 piscivorous developing an attack without “S-shape”. In this sequence, pikeperch move slowly
79 towards the prey and attack it with a fast tail movement projecting it rapidly forwards
80 (Turesson et al., 2002). Piscivorous behavior in a pikeperch population, can appears very
81 early for some individuals (11.0 ± 1.3 mm TL, 21 dph; Colchen et al., 2020a). It is exhibited by
82 all individuals four months post-hatching (70-80 mm; Persson and Brönmark, 2008). This
83 suggests that the precocious piscivores could become cannibals under farming conditions
84 because they have no other fish prey in their environment except their conspecifics.

85 In nature, pikeperch is an active predator during low light intensity (dawn and dusk;
86 Dörner et al., 2007) with a perfect visual acuity (Ali et al., 1977; Colchen et al., 2020b). The
87 transition to a piscivorous diet is governed by both anatomical modifications and the
88 availability of prey (Buijse and Houthuijzen, 1992; Claessen et al., 2002; Ribeiro et al., 2021).
89 Adult pikeperch prefer smaller prey than their body size (predator-prey length ratio: 0.23 on
90 average; Dörner et al., 2007) due to their anatomical (large gape) and digestive (presence of
91 enzymes) abilities. This prey/predator size ratio is the “predation window”, which could

92 fluctuate with available prey (Claessen et al., 2002) and, thus, impact the shift to the
93 piscivorous diet.

94 The main objective of this study was to test the possible link between cannibalism on
95 pikeperch and the onset of piscivorous behavior on a non-conspecific species, zebrafish
96 *Danio rerio*, known to display escape behavior and has a reactive distance for avoiding
97 perciform predators (Dill, 1974a; b). We hypothesized that a non-cannibal at a given time is
98 not able to display an efficient piscivorous behavior. This statement assumes that to be
99 cannibal, pikeperch larvae needs to be able to realize a full behavioral sequence to capture a
100 fish prey (describe in Colchen et al., 2019). We hypothesize that in larval and juvenile captive
101 populations of pikeperch, cannibalism could be based on the presence of early piscivorous
102 fish. As pikeperch is a strictly piscivorous fish, in a single-species farm conditions the onset of
103 piscivory means the onset of cannibalism, and consequently suggests that cannibalism is
104 unavoidable.

105

106 **2. Materials and Methods**

107 *2.1. Ethical note*

108 During all procedures, we took care to minimize handling and stress as much as
109 possible for the studied animals. All fish treatments and procedures used in this study were
110 in accordance with the guidelines of the Council of the European Union (2010/63/UE) and
111 approved by the French Animal Care Guidelines (Animal approval No. APAFIS#1813-
112 2015111618046759v2).

113

114 *2.2. Origin and larval rearing conditions*

115 Eggs of pikeperch came from two mature females fertilized by one male in a fish farm
116 (SARL Asialor, Pierrevillers, Moselle, France). After hatching (1st March, 2017), larvae were
117 reared at the Aquaculture Experimental Platform (AEP, registration number for animal
118 experimentation C54-547-18) belonging to the URAFPA lab located at the Faculty of Sciences
119 and Technologies of the University of Lorraine (Nancy-France) in a Recirculating Aquatic
120 System (RAS). They were reared until 14 dph in 700L tanks. Artificial lighting (50 lx at the
121 water surface) followed a 12L/12D cycle with light on from 08.00 am to 8.00 pm with 30 min
122 simulation of dawn and dusk. The water was maintained at 16°C during the first two days
123 and increased by 1°C per day until reaching 20°C. At 14 dph, larvae were transferred from
124 700L tanks to four 300L independent incubators at 20.0°C ± 0.0°C each containing nine
125 enclosures (38 cm x 7 cm x 11 cm of deep each enclosure with 2.5 L of water). Three
126 enclosures per incubators contained 100 pikeperch larvae for cannibalism observations and
127 the six other ones contained cages (four cages per enclosure) for cannibals and non-
128 cannibals larval rearing. Each incubator was a recirculating water system (110 x 64 x 186 cm)
129 with a flow rate of 4 m³.h⁻¹ and water was UV sterilized (for more information about the
130 experimental facilities, see Réalis-Doyelle et al., 2016; 2022). Incubators were cleaned every
131 morning. Water parameters (mean ± standard deviation, SD) were measured once a week:
132 dissolved oxygen (8.28 ± 0.98 mg.L⁻¹); pH (7.83 ± 0.05); ammonia (NH₄⁺, 0 mg.L⁻¹); nitrite
133 (NO₂⁻, 0.32 ± 0.14 mg.L⁻¹). Larvae were fed live prey and commercial inert feed seven times
134 per day as follows: *Artemia* nauplii (550-600 µm, Catvis, Hertogenbosch, Pays-Bas) from 4 to
135 18 dph, then, Larviva PROWEAN 100, 300, 500, 700 µm (BioMar®, France) until 35 dph and
136 INICIOplus 0.8 mm (BioMar®, France) after 35 dph.

137

138 2.3. Selection of cannibals and non-cannibals

139 For this experiment, predatory behaviors of cannibals were compared to those of
140 non-cannibals. To obtain cannibals, larvae were reared in three enclosures located in each
141 incubator and were identified during observation sessions from above view by focal
142 sampling of five minutes each hour during the light period (a density of 100 larvae per
143 enclosure was used). Cannibals were identified by the presence of a prey in its mouth. If
144 cannibalism was observed, the cannibal, which tends to swim at the surface of the tank with
145 the prey in its mouth, was captured and transferred with a fine mesh dip net into an
146 individual cage (9 cm diameter x 10 cm of height, with 7 cm of water depth and 0.45 L of
147 water) set up in the other six enclosures of each incubator. To obtain non-cannibals, other
148 pikeperch larvae were reared in a cage (9 cm diameter x 10 cm of height, with 7 cm of water
149 depth and 0.45 L of water) in groups of four individuals, and mortality was observed every
150 day for 17 days. If all four larvae were alive the day of the test, they were all considered as
151 non-cannibal for the test. As a total, 96 cages with four larvae were used at the beginning of
152 the experiment to obtain non-cannibals.

153 We obtained 59 cannibals and 59 non-cannibals aged of 17 to 34 dph (between L5
154 and L6 stages; Ott et al., 2012). Two cannibals died during the isolation period, one cannibal
155 and one non-cannibal died between the two tests and four videos could technically not be
156 used. Consequently, eight pairs of cannibals and non-cannibals were deleted from the
157 analysis. In total, behaviors of 51 cannibals (mean of TL \pm SD: 19.4 \pm 4.7 mm) and 51 non-
158 cannibals (mean of TL \pm SD: 17.1 \pm 2.6 mm) were analyzed. Total length was significantly
159 different between cannibals and non-cannibals (U = 1647; p = 0.006). The behavioral tests
160 started when the first case of cannibalism was observed.

161

162 *2.4. Behavioral tests*

163 Two behavioral tests were conducted: a predation test and a cannibalism test. For
164 each trial, one cannibal and one non-cannibal of the same age were each isolated in a cage
165 and not fed during a 24-hour period before the test, then introduced into two rectangular
166 arenas (20 x 7 x 4 cm with 2 cm of water height, 280 mL): one arena for cannibal and one
167 arena for non-cannibal. These arenas were identical and both divided in two equal zones by
168 an opaque removable divider. Cannibal and non-cannibal were introduced, respectively, in
169 the first zone of each rectangular arena and acclimatized for 30 minutes. The type of test
170 depended of individuals introduced in the second zone at the same time of tested larvae.

171 For predation test, three non-conspecific prey (zebrafish larvae, mean of TL \pm SD: 4.0
172 \pm 0.8 mm) were introduced in the second zone of each arena and let for the same
173 acclimatization period than tested larvae. After 30 minutes, the divider of each arena was
174 removed allowing the tested larva (cannibal or non-cannibal) and the three zebrafish of each
175 arena to explore it. For cannibalism test, three conspecifics, which were non-cannibals,
176 (pikeperch larvae, mean of TL \pm SD: 14.3 \pm 2.3 mm) were introduced in the second zone of
177 each rectangular arena and acclimatized for 30 minutes. After 30 minutes, the divider of
178 each arena was removed allowing the tested larva (cannibal or non-cannibal) and the three
179 conspecifics of each arena to explore it.

180 For each tested larva, the test order was randomized. After the first test (predation
181 or cannibalism test), pikeperch, cannibal and non-cannibal, were put back in their own cage
182 and the second test (predation or cannibalism in function of the first conducted test) was
183 conducted 24 hours later. The fish behaviors were video recorded with two camcorders
184 (Sony, Handycam, DCR-SR72E) positioned 80 cm above each arena for 20 minutes. The
185 acclimatization period was not considered. For each test, measured parameters concerning
186 the predator or the cannibal were: the number of orientations, approaches, attacks (with "S-

187 shape” and without “S-shape”), pursuits and captures (for the definitions see **Table 1**). The
188 cannibal or non-cannibal performance was calculated as follows: number of
189 captures/number of attacks.

190

191 *2.5. Statistical analyses*

192 The total length of larvae of each group was statistically compared with a Mann-
193 Whitney U-test, and the influence of the test order (cannibalism test then predation test and
194 vice versa) was tested on all behavioral parameters by Generalized Linear Model, (package
195 ‘glmmADMB’; Zuur et al., 2009).

196 Data did not fit with normality (Shapiro-Wild test, shapiro.test (R Core Team, 2017))
197 and variance homogeneity (leveneTest package “car” (Fox and Weisberg, 2016)) for
198 behavioral parameters: the number of orientations, approaches, attacks with “S-shape” or
199 without “S-shape”, captures, pursuits and performance. Comparisons of behavioral
200 responses between cannibals and non-cannibals were analyzed with Mann-Whitney U-test
201 for independent data and comparisons between the cannibalism and predation test were
202 analyzed with Wilcoxon test for dependent data. To compare the number of attacks with “S-
203 shape” with attacks without “S-shape”, a model based on rank transformation was used with
204 a tested interaction Tests*Types where ‘Tests’ corresponding to cannibalism and predation
205 tests and ‘Types’ corresponding to attacks with or without the “S-shape” behavior (packages
206 “ordinal” (Christensen, 2015) and “ART” (Villacorta, 2015)). When a significant interaction
207 was found, a pairwise comparison using Tukey-Kramer-Nemenyi with Tukey-Dist
208 approximation was performed (package “PMCMRplus” (Pohlert, 2022)). All statistical
209 analyses were performed using the free software Rstudio (version 2021.09.2). Results were
210 presented by mean \pm SD. The level of significance used in all tests was $p_{\text{adjust}} < 0.05$.

211

212 3. Results

213 3.1. Comparisons between cannibals and non-cannibals

214 First of all, behavioral responses were independent of the test sequence (**Table 2**).

215 In the predation test on zebrafish, behaviors differed significantly between cannibals
216 and non-cannibals. Cannibals compared to non-cannibals displayed fewer orientations ($14 \pm$
217 11 vs. 30 ± 20 ; $W = 640$; $p_{\text{adjust}} < 0.0001$; **Fig. 1A; Table 3**), less approaches (9 ± 7 vs. 17 ± 12 ;
218 $W = 704$; $p_{\text{adjust}} = 0.0002$; **Fig. 1B; Table 3**), less attacks (9 ± 8 vs. 19 ± 15 ; $W = 758$; $p_{\text{adjust}} =$
219 0.0008 ; **Fig. 1C; Table 3**), but they caught more zebrafish (2 ± 1 vs. 0 ± 1 ; $W = 2004$; $p_{\text{adjust}} <$
220 0.0001 ; **Fig. 1D; Table 3**). There was no difference for the number of pursuits between
221 cannibals and non-cannibals ($W = 1027$; $p_{\text{adjust}} = 0.1$; **Fig.1E; Table 3**). Cannibals were
222 significantly more efficient than non-cannibals (0.31 ± 0.28 vs. 0.05 ± 0.15 ; $W = 1872$; $p_{\text{adjust}} <$
223 0.0001 ; **Fig. 1F; Table 3**).

224 In the cannibalism test, the number of approaches (23 ± 21 vs. 17 ± 21 ; $W = 1622$;
225 $p_{\text{adjust}} = 0.03$; **Fig. 1B**), attacks (4 ± 4 vs. 3 ± 6 ; $W = 1660$; $p_{\text{adjust}} = 0.01$; **Fig. 1C**), captures (1 ± 1
226 vs. 0 ± 0 ; $W = 1683$; $p_{\text{adjust}} = 0.0001$; **Fig. 1D**) and the performance (0.21 ± 0.29 vs. 0.02 ± 0.08 ;
227 $W = 700$; $p_{\text{adjust}} = 0.0006$; **Fig. 1F**) were higher among cannibals than non-cannibals. Number
228 of orientations and pursuits did not differ significantly between the two groups ($W = 1602$;
229 $p_{\text{adjust}} = 0.05$; **Fig. 1A; Table 3** and $W = 1418$; $p_{\text{adjust}} = 0.2$; **Fig. 1E; Table 3**, respectively).

230

231 3.2. Comparisons of behavioral performances in predation and cannibalism tests

232 Cannibals performed more orientations and approaches to congeners in the
233 cannibalism test than to prey in the predation test (**Table 3**). They realized also more attacks,
234 captures and pursuits in the predation than in the cannibalism test (**Table 3**). Cannibals

235 exhibited better performance in the predation than in the cannibalism test (**Table 3**). Non-
236 cannibals realized more attacks, captures and pursuits in the predation test than in the
237 cannibalism test (**Table 3**). For non-cannibals, all others analyzed behaviors did not differ
238 (**Table 3**).

239 Cannibals displayed more attacks with “S-shape” in the predation test (mean \pm SD: 6
240 \pm 6) than in cannibalism test (mean \pm SD: 2 \pm 3; $W = 120$; $p_{\text{adjust}} < 0.0001$) and also more
241 attacks without “S-shape” in predation test (mean \pm SD: 3 \pm 5) than in the cannibalism test
242 (mean \pm SD: 2 \pm 2; $W = 178$; $p_{\text{adjust}} = 0.002$). Non-cannibals displayed more attacks with “S-
243 shape” in the predation test (mean \pm SD: 12 \pm 9) than in the cannibalism test (mean \pm SD: 2 \pm
244 6; $W = 57.6$; $p_{\text{adjust}} < 0.0001$) and also more attacks without “S-shape” in the predation test
245 (mean \pm SD: 7 \pm 10) than in the cannibalism test (mean \pm SD: 1 \pm 2; $W = 56$; $p_{\text{adjust}} < 0.0001$).

246

247 *3.3. Comparisons between both types of attacks in each test for cannibals and non-*
248 *cannibals*

249 For cannibals in the cannibalism test, there was no difference between the number
250 of attacks with “S-shape” and without “S-shape” ($q = 2.1$; $p_{\text{adjust}} = 0.12$; **Fig. 2A**). However, in
251 the predation test they significantly displayed more attacks with “S-shape” than without ($q =$
252 4.5 ; $p_{\text{adjust}} = 0.001$; **Fig. 2A**). Non-cannibals displayed significantly more attacks than
253 cannibals using “S-shape” in the two tests (cannibalism test: $q = 3.4$; $p_{\text{adjust}} = 0.02$; predation
254 test: $q = 4.8$; $p_{\text{adjust}} = 0.0007$; **Fig. 2B**).

255

256 **4. Discussion**

257 This study highlights a direct link between the onset of piscivory and cannibalism in
258 pikeperch larvae. Cannibals capture both prey and conspecifics with higher performance

259 whereas non-cannibals catch little prey with lower efficiency and no conspecifics (**Fig. 1**).
260 Cannibals were better predators in the predation test than in the cannibalism test (**Table 3**).
261 Overall, cannibals were better predators than non-cannibals whatever the prey tested,
262 developing predatory behavioral items for piscivory before their conspecifics. These results
263 contribute to the understanding of cannibalism, a major bottleneck for rearing of pikeperch,
264 a commercially important species.

265 The foraging bouts of piscivorous fish correspond to a similar sequence across
266 species: prey search, recognition and localization (defined by orientation and approach in
267 the present study), attack, capture and ingestion (Holling, 1965; Endler, 1991). The present
268 study demonstrates this sequence in pikeperch larvae, which is known to be a prerequisite
269 for predator and particularly for fish larvae (e.g. common carp *Cyprinus carpio* and Northern
270 pike *Esox Lucius* (Drost, 1987)). Behaviors implied in this behavioral sequence were few
271 studied in a piscivory context (Howick and O'Brien, 1983; Turesson and Brönmark, 2004). In
272 our study, we showed that non-cannibals did not fully control the behavioral sequence for
273 fish capture contrary to cannibals of the same age. They displayed lot of orientations and
274 approaches without attacking or when they attacked, they do not succeed to capture
275 potential fish prey. Non-cannibal larvae were still in a learning process, particularly the
276 acquisition of a good attack behavior. Such an optimal attack distance to succeed a capture
277 was already demonstrated in chain pickerel *Esox niger* (Rand and Lauder, 1981) and
278 pikeperch (Colchen et al., 2020a). This estimation of the best attack distance, correlated with
279 fish age and development, is not an innate process and predators try several times to attack
280 a prey before being able to control it (Colchen et al., 2020a).

281 Piscivorous fishes develop different tactics for capturing prey, such as ambushing,
282 stalking and chasing (Takeuchi, 2009), in parallel with morphological traits, such as oral jaw

283 teeth, which are crucial for both capture and handling prey in 12 fish species (Muruga et al.,
284 2022). These diverse predator tactics could be related to the wide range of behavioral
285 adaptations for avoiding capture, among which escape behavior in roach *Rutilus rutilus*
286 (Ranåker et al., 2012), freezing in glow light tetras *Hemigrammus erythrozonus* (Brown and
287 Dreier, 2002) or attacks to the predator as mobbing in French grunt *Haemulon flavolineatum*
288 (Hein, 1996). In our tests, zebrafish and pikeperch larvae have different abilities to defend
289 themselves against a predator. Zebrafish use the escape behavior and their reactive distance
290 for avoiding largemouth bass *Micropterus salmoides*, a real predator, and artificial predators
291 (model predator and a film of an approaching object) depends on both the size and velocity
292 of the predator and not on the distance from the predator or the predator itself (Dill, 1974a;
293 b). Here, pikeperch larvae used as prey were between L5 and L6 stages, with spiny rays on
294 dorsal fins, and a complete development of fins and teeth (Ott et al., 2012) whereas
295 zebrafish larvae had neither developed dorsal fin nor teeth at 4 mm (Parichy et al., 2009),
296 which could partly explain the difference between predation and cannibalism tests. Piscivory
297 was not only reliant on predator selectivity, but also, how prey influence their relative
298 catchability, implying for predator modifications of their capture strategy (Mihalitsis et al.,
299 2021). For a cannibal, it is arguably riskier to attack a conspecific than a non-conspecific prey
300 with less risky structures (fin rays and body spines), such as zebrafish. Cannibals displayed lot
301 of orientations and approaches towards conspecifics, as non-cannibals in predation test
302 towards prey, before attacking. Even if the behavioral sequence to attack and capture a
303 conspecific or a non-conspecific prey seems to be similar, there are quantitative differences
304 in displayed behaviors, with more orientations, approaches and attacks in the cannibalism
305 test, showing that cannibals are probably able to identify a conspecific as potentially
306 dangerous. Consequently, it seems that cannibals choose an attack tactic with less risks

307 against a conspecific. To attack non-conspecific prey, cannibals used “S-shape”. However, to
308 attack conspecific prey, they used “S-shape” but also no “S-shape” tactic. It highlights that
309 cannibals try to shift to a better capture method but without a complete control of it. The
310 attack with “S-shape”, imposing a pause before attack, is characteristic for some fish larvae
311 attacking zooplankton and invertebrates and represented an important part of type of
312 attack: 50 % in pikeperch larvae (34.6 ± 9.4 mm TL; Colchen et al., 2019; 2020a) and 31.25 %
313 in European whitefish *Coregonus lavaretus* larvae (15.5 ± 0.3 mm TL; Mahjoub et al., 2008).
314 In diet transition, there is a gap from zooplankton and invertebrates to fish (piscivory)
315 feeding implying behavioral changes (Sánchez-Hernández et al., 2019) already described in
316 pikeperch or in walleye (Persson and Brönmark, 2002; Graeb et al., 2005). When pikeperch
317 shift to a piscivorous diet, they need to change their capture method, by switching to an
318 attack without “S-shape” marked by a tail propulsion (appearing between 35 and 40 dph,
319 32.7 ± 4.2 mm TL, unpublished data), corresponding to the attack without “S-shape”
320 (Sullivan and Atchinson, 1978). Hunting behavioral transition was mentioned in some marine
321 fish larvae where the time spent poised in a striking posture (*i.e.* “S-shape” position) is much
322 longer during first feeding than in later larval stages, and gradually the strike becomes
323 integrated with swimming movements (as attack without “S-shape”) (Hunter, 1980). A
324 poised striking posture seems to be a common tendency in young larvae and declines as
325 larvae grow up suggesting that it could be an adaptation to low feeding success (Hunter,
326 1980). On one hand, the absence of difference between both types of attacks for cannibals
327 showed that they were in progress to change their attack method (from “S-shape” to non “S-
328 shape” tactic), due to the probably decreasing success to catch prey with the first tactic. On
329 the other hand, by using “S-shape” behavior to attack indistinctly conspecific or non-
330 conspecific, non-cannibals showed that they did not acquire the good attack tactic to

331 capture fish prey. Nevertheless, the success of capture for non-cannibals in predation test
332 with “S-shape” attacks, even if prey were fish, could be explained by the small size of
333 zebrafish larvae. Indeed, handling times to capture zebrafish could consider as negligible like
334 for zooplankton (Hunter, 1980). Furthermore, there is a favorite size range of prey for
335 predatory fishes. In function of species, maximal prey consumed size observed was between
336 35 and 78 % of predatory size (Persson et al., 2000; Ribeiro and Qin, 2013). It was previously
337 suggested that 50 % of size difference could be the threshold to initiate intracohort
338 cannibalism in Barramundi *Lates calcarifer* (Ribeiro and Qin, 2013). In pikeperch larvae, this
339 ratio was not known to us but in the present study the ratio between prey size and predator
340 size was 0.8 ± 0.1 for cannibals and non-cannibals in cannibalism test and 0.2 ± 0.0 in
341 predation test. We can conclude that cannibals and non-cannibals had the same size abilities
342 to capture congeners and zebrafish larvae. However, for larger fish prey, as conspecifics, it
343 requires a new set of motor patterns associated with grasping prey and presence of
344 anatomical characteristics (e.g.: teeth or mouth gape; Galarowicz and Wahl, 2005).

345 Therefore, under monospecific farming conditions, piscivorous feeding was reported
346 to start very early in development of several fish species like perch (10-13 mm; Hunter,
347 1980; Brabrand, 1995) or walleye (30 mm; Graeb et al., 2005). The present study
348 demonstrated that cannibalism is linked to the onset of available piscivory and not to the
349 onset of predatory behavior that was already controlled when larvae catch zooplankton.
350 Previously, it was demonstrated that very early in the development, pikeperch larvae could
351 capture *Artemia* nauplii but are not able to capture fish prey (Colchen et al., 2020a). Here,
352 we highlighted that all fish able to capture fish prey are potentially cannibals, or at least, that
353 early piscivorous fish have a high probability to become cannibals.

354

355 5. Conclusions

356 To conclude, our study showed that cannibalism is linked to the onset of piscivory
357 and to the acquisition of the behavioral sequence of predation. This statement could explain
358 that when cannibalism is established in a pikeperch population in farming conditions, it is
359 probably irreversible (Baras et Jobling, 2002; Steenfeldt et al., 2015). Are there solutions to
360 avoid cannibalism in pikeperch larvae? Cannibals showed a faster development and are
361 rapidly larger than their conspecifics. A first solution is to practice the sorting of the
362 individuals and to eliminate the largest ones (Naumowicz et al., 2017). This method, helping
363 to reach higher survival rates, mainly due to lower losses caused by cannibalism
364 (Szczepkowski et al., 2011), needs to be repeated frequently (every 10 days for 0.7-8.2 g fish
365 and afterwards at 21 days intervals (Policar et al., 2016)) because larger fish appear quickly
366 in the population. The second method consists to eliminate the fish that exhibit a
367 cannibalistic behavior. This method could be based on a genetic process with individual
368 selection based on later ichthyophagy (*i.e.* Norton et al., 2011; Jensen, 2015).

369 In order to select fishes whose behaviors make them more suitable for aquaculture
370 production, this study could be useful to investigate in wild and domestic strains behavioral
371 parameters implied in cannibalism (*i.e.* aggressiveness; Toomey et al., 2019) to establish an
372 early diagnosis of the cannibal potential of a population. This investigation is supported by
373 that cannibalistic behavior was already showed as genetically-based to a great extent in
374 hybrid catfish *Silurus meridionalis-asotus* (Yang et al., 2015) and that aggressiveness can be
375 attributed to the combination or interaction of genotypic variation and environmental
376 difference in sticklebacks *Gasterosteus aculeatus* (Bakker's, 1986). In pikeperch, producing a
377 population with the lowest cannibal potential is the ultimate goal. However, there is a huge
378 variability in behavior at the intraspecific level (Toomey et al., 2019; 2020).

379

380 **6. Acknowledgements**

381 We would like to thank SARL Asialor, and particularly M. Alix, J.B. Muliloto and D. Źarski for
382 supplying pikeperch larvae and A. Forfert and L. Lorrain for their help to analyze some
383 videos.

384

385 **7. References**

386 Ali, M.A., Ryder, R.A., Anctil, M., 1977. Photoreceptors and visual pigments as related to
387 behavioral responses and preferred habitats of perches (*Perca* spp.) and pikeperches
388 (*Stizostedion* spp.). Journal of the Fisheries Research Board of Canada 34, 1475–1480.
389 DOI: 10.1139/f77-212

390 Bakker, T.C., 1986. Aggressiveness in sticklebacks (*Gasterosteus aculeatus* L.): a behaviour-
391 genetic study. Behaviour, 98(1-4), 1-144.

392 Baras, E., Jobling, M., 2002. Dynamics of intracohort cannibalism in cultured fish.
393 Aquaculture research 33, 461–479. DOI: 10.1046/j.1365-2109.2002.00732.x

394 Bell, A.M., & Sih, A., 2007. Exposure to predation generates personality in threespined
395 sticklebacks (*Gasterosteus aculeatus*). Ecology letters, 10(9), 828-834. DOI :
396 10.1111/j.1461-0248.2007.01081.x

397 Brabrand, A., 1995. Intra-cohort cannibalism among larval stages of perch (*Perca*
398 *fluviatilis*). Ecology of Freshwater Fish, 4(2), 70-76. DOI: 10.1111/j.1600-
399 0633.1995.tb00119.x

400 Brown, G.E., Dreier, V. M., 2002. Predator inspection behavior and attack cone avoidance in
401 a characin fish: the effects of predator diet and prey experience. Animal
402 Behavior, 63(6), 1175-1181. DOI: 10.1006/anbe.2002.3024

403 Buijse, A.D., Houthuijzen, R.P., 1992. Piscivory, growth, and size-selective mortality of age 0
404 pikeperch (*Stizostedion lucioperca*). Canadian Journal of Fisheries and Aquatic Sciences
405 49, 894–902. DOI:10.1139/f92-100

406 Christensen, R.H.B., 2015. ordinal - Regression Models for Ordinal Data. R package version
407 2015.6-28. <http://www.cran.r-project.org/package=ordinal/>.

408 Claessen, D., Van Oss, C., de Roos, A.M., Persson, L., 2002. The impact of size-dependent
409 predation on population dynamics and individual life history. Ecology 83, 1660–1675.
410 DOI: 10.1890/0012-9658(2002)083[1660:TIOSDP]2.0.CO;2

411 Colchen, T., Teletchea, F., Fontaine, P., Pasquet, A., 2017. Temperature modifies activity,
412 inter-individual relationships and group structure in fish. Current Zoology 163, 175–
413 183. DOI:10.1093/cz/zow048.

414 Colchen, T., Fontaine, P., Ledoré, Y., Teletchea, F., Pasquet, A., 2019. Intra-cohort
415 cannibalism in early life stages of pikeperch. Aquaculture Research 50, 915-924. DOI:
416 10.1111/are.13966.

417 Colchen, T., Dias, A., Gisbert, E., Teletchea, F., Fontaine, P., Pasquet, A., 2020a. The onset of
418 piscivory in fish: behavioral and physiological correlations. Journal of Fish Biology
419 96:1463-1474. DOI: 10.1111/jfb.14322.

420 Colchen, T., Gisbert, E., Ledoré, Y., Teletchea, F., Fontaine, P., Pasquet, A., 2020b. Is a
421 cannibal different from its conspecifics? A behavioral, morphological, muscular and
422 retinal structure study with pikeperch juveniles under farming conditions. Applied
423 Animal Behavior Science 224. DOI: 10.1016/j.applanim.2020.104947.

424 Cuff, W.R., 1980. Behavioral aspects of cannibalism in larval walleye, *Stizostedion*
425 *vitreum*. Canadian Journal of Zoology, 58(8), 1504-1507. DOI: 10.1139/z80-207

426 Curio, E., 2012. The ethology of predation (Vol. 7). Springer Science & Business Media.

427 Dill, L.M., 1974a. The escape response of the zebra danio (*Brachydanio rerio*) I. The stimulus
428 for escape. *Animal Behavior*, 22(3), 711-722. DOI: 10.1016/S0003-3472(74)80022-9

429 Dill, L.M., 1974b. The escape response of the zebra danio (*Brachydanio rerio*) II. The effect of
430 experience. *Animal Behaviour*, 22(3), 723-730. DOI: 10.1016/S0003-3472(74)80023-0

431 Dörner, H., Hülsmann, S., Hölker, F., Skov, C., Wagner, A., 2007. Size-dependent predator-
432 prey relationships between pikeperch and their prey fish. *Ecology of Freshwater Fish*
433 16, 307–314. DOI:10.1111/j.1600-0633.2006.00223.x

434 Drost, M.R., 1987. Relation between aiming and catch success in larval fishes. *Canadian*
435 *Journal of Fisheries and Aquatic Sciences*, 44(2), 304-315. DOI: 10.1139/f87-039

436 Endler, J.A., 1991. Variation in the appearance of guppy color patterns to guppies and their
437 predators under different visual conditions. *Vision research* 31, 587–608. DOI:
438 10.1016/0042-6989(91)90109-1

439 FAO. *Sander lucioperca* (Linnaeus, 1758). Fisheries and Aquaculture Division, Rome. 2021.
440 Available online: <https://www.fao.org/fishery/en/aqspecies/3098> (accessed on 3
441 February 2023).

442 Fox, J., Weisberg, S., 2016. Package ‘car’ Companion to Applied Regression. R Package
443 version, 2–1. <https://cran.r-project.org/web/packages/car/car.pdf>.

444 Galarowicz, T.L., Wahl, D.H., 2005. Foraging by a young-of-the-year piscivore: the role of
445 predator size, prey type, and density. *Canadian Journal of Fisheries and Aquatic*
446 *Sciences*, 62(10), 2330-2342. DOI: 10.1139/f05-148

447 Graeb, B.D., Galarowicz, T., Wahl, D.H., Dettmers, J.M., Simpson, M.J., 2005. Foraging
448 behavior, morphology, and life history variation determine the ontogeny of piscivory in
449 two closely related predators. *Canadian Journal of Fisheries and Aquatic*
450 *Sciences*, 62(9), 2010-2020. DOI: 10.1139/f05-112

451 Holling, C.S., 1965. The functional response of predators to prey density and its role in
452 mimicry and population regulation. The Memoirs of the Entomological Society of
453 Canada 97, 5–60. DOI:10.4039/entm9745fv

454 Houde, E.D., 2001. Fish larvae. In: *Encyclopedia of Ocean Sciences* (Ed. Steele, J.H., Turekian,
455 K.K. and Thorpe, S.A.). pp. 928-938. London: Academic Press.

456 Howick, G.L., O'Brien, W.J., 1983. Piscivorous feeding behavior of largemouth bass: an
457 experimental analysis. *Transactions of the American Fisheries Society*, 112(4), 508-516.
458 DOI: 10.1577/1548-8659(1983)112%3c508:PFBOLB%3e2.0.CO;2

459 Hunter, J.R., 1980. The feeding behavior and ecology of marine fish larvae. In *Fish behavior*
460 *and its use in the capture and culture of fishes*, ICLARM Conference Proceedings (pp.
461 287-330).

462 Hein, R.G., 1996. Mobbing behavior in juvenile French grunts (*Haemulon*
463 *flavolineatum*). *Copeia*, 1996(4), 989-991. DOI: 10.2307/1447662

464 Jensen, P., 2015. Adding 'epi-'to behavior genetics: implications for animal
465 domestication. *Journal of Experimental Biology*, 218(1), 32-40. DOI:
466 10.1242/jeb.106799

467 Kaji, T., Kodama, M., Arai, H., Tagawa, M., Tanaka, M., 2002. Precocious development of the
468 digestive system in relation to early appearance of piscivory in striped bonito *Sarda*
469 *orientalis* larvae. *Fisheries science*, 68(6), 1212-1218. DOI: 10.1046/j.1444-
470 2906.2002.00557.x

471 Kestemont, P., Jourdan, S., Houbart, M., Mélard, C., Paspatis, M., Fontaine, P., Cuvier, A.,
472 Kentouri, M., Baras, E., 2003. Size heterogeneity, cannibalism and competition in
473 cultured predatory fish larvae: biotic and abiotic influences. *Aquaculture* 227, 333–
474 356. doi:10.1016/S0044-8486(03)00513-1

475 Kohlmeier, C., Ebenhöf, W., 1995. The stabilizing role of cannibalism in a predator-prey
476 system. *Bulletin of Mathematical Biology* 57, 401–411. DOI: 10.1007/BF02460632

477 Kucharczyk, D., Kestemont, P., Mamcarz, A., 2007. Artificial reproduction of pikeperch.
478 Mercurius, Olsztyn, Poland.

479 Krebs, J.R., Davies, N.B., 2009. *Behavioral ecology: an evolutionary approach*. John Wiley &
480 Sons.

481 Lappalainen, J., Dorner, H., Wysujack, K., 2003. Reproduction biology of pikeperch (*Sander*
482 *lucioperca* (L.)) - a review. *Ecology of Freshwater Fish* 12(2), 95–106. DOI:
483 10.1034/j.1600-0633.2003.00005.x

484 MacArthur, R.H., Pianka, E., 1966. On optimal use of patchy environment. *The American*
485 *Naturalist* 100, 603–609. DOI: 10.1086/282454

486 Mahjoub, M. S., Anneville, O., Molinero, J. C., Souissi, S., Hwang, J. S., 2008. Feeding
487 mechanism and capture success of European whitefish (*Coregonus lavaretus* L.)
488 larvae. *Knowledge and Management of Aquatic Ecosystems*, (388), 05. DOI:
489 10.1051/kmae:2008008

490 Mihalitsis, M., Hemingson, C. R., Goatley, C. H., Bellwood, D. R., 2021. The role of fishes as
491 food: A functional perspective on predator–prey interactions. *Functional Ecology* 35(5),
492 1109-1119. DOI: 10.1111/1365-2435.13779

493 Muruga, P., Bellwood, D. R., Mihalitsis, M., 2022. Forensic odontology: Assessing bite
494 wounds to determine the role of teeth in piscivorous fishes. *Integrative Organismal*
495 *Biology*, 4(1), obac011. DOI: 10.1093/iob/obac011

496 Naumowicz, K., Pajdak, J., Terech-Majewska, E., Szarek, J., 2017. Intracohort cannibalism and
497 methods for its mitigation in cultured freshwater fish. *Reviews in Fish Biology and*
498 *Fisheries*, 27(1), 193-208. DOI: 10.1007/s11160-017-9465-2

499 Norton, W.H., Stumpenhorst, K., Faus-Kessler, T., Folchert, A., Rohner, N., Harris, M.P.,
500 Callebert J., Bally-Cuif, L., 2011. Modulation of Fgfr1a signalling in zebrafish reveals a
501 genetic basis for the aggression–boldness syndrome. *Journal of Neuroscience*, 31(39),
502 13796-13807. DOI: 10.1523/JNEUROSCI.2892-11.2011

503 Ott, A., Löffler, J., Ahnelt, H., Keckeis, H., 2012. Early development of the postcranial
504 skeleton of the pikeperch *Sander lucioperca* (Teleostei: Percidae) relating to
505 developmental stages and growth. *Journal of Morphology* 273, 894–908.
506 DOI:10.1002/jmor.20029

507 Parichy, D. M., Elizondo, M. R., Mills, M. G., Gordon, T. N., Engeszer, R. E., 2009. Normal
508 table of postembryonic zebrafish development: staging by externally visible anatomy
509 of the living fish. *Developmental dynamics*, 238(12), 2975-3015. DOI:
510 10.1002/dvdy.22113

511 Persson, L., Byström, P., & Wahlström, E., 2000. Cannibalism and competition in Eurasian
512 perch: population dynamics of an ontogenetic omnivore. *Ecology*, 81(4), 1058-1071.
513 DOI: 10.1890/0012-9658(2000)081[1058:CACIEP]2.0.CO;2

514 Persson, A., Brönmark, C., 2002. Foraging capacity and resource synchronization in an
515 ontogenetic diet switcher, pikeperch (*Stizostedion lucioperca*). *Ecology*, 83(11), 3014-
516 3022. DOI: 10.1890/0012-9658(2002)083[3014:FCARSI]2.0.CO;2

517 Persson, A., Brönmark, C., 2008. Pikeperch *Sander lucioperca* trapped between niches:
518 foraging performance and prey selection in a piscivore on a planktivore diet. *Journal of*
519 *Fish Biology* 73, 793–808. DOI:10.1111/j.1095-8649.2008.01956.x

520 Pohlert, T., 2022. PMCMRplus: Calculate Pairwise Multiple Comparisons of Mean Rank Sums
521 Extended. R package version 1.9.6. [https://cran.r-](https://cran.r-project.org/web/packages/PMCMRplus/index.html)
522 [project.org/web/packages/PMCMRplus/index.html](https://cran.r-project.org/web/packages/PMCMRplus/index.html)

523 Policar, T., Křišťan, J., Blecha, M., & Vaniš, J., 2016. Adaptation and culture of pikeperch
524 (*Sander lucioperca* L.) juveniles in recirculating aquaculture system (RAS). University of
525 South Bohemia in České Budějovice, Faculty of Fisheries and Protection of Waters.

526 Policar, T., Schaefer, F. J., Panana, E., Meyer, S., Teerlinck, S., Toner, D., Źarski, D., 2019.
527 Recent progress in European percid fish culture pro- duction technology-tackling
528 bottlenecks. *Aquaculture International* 27 (5), 1151–1174. DOI: 10.1007/s10499-019-
529 00433-y

530 Polis, G.A., 1981. The evolution and dynamics of intraspecific predation. *Annual Review of*
531 *Ecology and Systematics* 12, 225–251. <http://www.jstor.org/stable/2097111>.

532 R Core Team, 2017. R: a Language and Environment for Statistical Computing, version 3.2.4.
533 R foundation for statistical computing, Vienna, Austria.

534 Ranåker, L., Jönsson, M., Nilsson, P. A., Brönmark, C., 2012. Effects of brown and turbid
535 water on piscivore–prey fish interactions along a visibility gradient. *Freshwater*
536 *biology*, 57(9), 1761-1768. DOI: 10.1111/j.1365-2427.2012.02836.x

537 Ranåker, L., Persson, J., Jönsson, M., Nilsson, P.A., Brönmark, C., 2014. Piscivore-Prey Fish
538 Interactions: Mechanisms behind Diurnal Patterns in Prey Selectivity in Brown and
539 Clear Water. *PLoS ONE* 9, e102002. DOI:10.1371/journal.pone.0102002

540 Rand, D. M., Lauder, G. V. 1981. Prey capture in the chain pickerel, *Esox niger*: correlations
541 between feeding and locomotor behavior. *Canadian Journal of Zoology*, 59(6), 1072-
542 1078. DOI: 10.1139/z81-149

543 Réalis-Doyelle, E., Pasquet, A., De Charleroy, D., Fontaine, P., Teletchea, F., 2016. Strong
544 effects of temperature on the early life stages of a cold stenothermal fish species,
545 brown trout (*Salmo trutta* L.). *PLoS One*, 11(5), e0155487. DOI:
546 10.1371/journal.pone.0155487

547 Réalis-Doyelle, E., Pasquet, A., Fontaine, P., Teletchea, F., 2022. Effects of temperature on
548 the survival and development of the early life stages of northern pike (*Esox*
549 *lucius*). Knowledge & Management of Aquatic Ecosystems, (423), 10. DOI:
550 10.1051/kmae/2022007

551 Ribeiro, F.F., & Qin, J.G., 2013. Modelling size-dependent cannibalism in barramundi *Lates*
552 *calcarifer*: cannibalistic polyphenism and its implication to aquaculture. PloS
553 one, 8(12), e82488. DOI : 10.1371/journal.pone.0082488

554 Ribeiro, D., Gkenas, C., Gago, J., Ribeiro, F., 2021. Variation in Diet Patterns of the Invasive
555 Top Predator *Sander lucioperca* (Linnaeus, 1758) across Portuguese
556 Basins. Water 13(15), 2053. DOI: 10.3390/w13152053

557 Sánchez-Hernández, J., Nunn, A. D., Adams, C. E., Amundsen, P. A., 2019. Causes and
558 consequences of ontogenetic dietary shifts: a global synthesis using fish
559 models. Biological Reviews, 94(2), 539-554. DOI: 10.1111/brv.12468

560 Steinfeldt, S.J., Lund, I., 2008. Development of methods of production for intensive rearing
561 of pikeperch juveniles. (No. 199). DTU Aqua Research Report, University of Denmark,
562 Denmark.

563 Steinfeldt, S., Lund, I., Höglund, E., 2011. Is batch variability in hatching time related to size
564 heterogeneity and cannibalism in pikeperch (*Sander lucioperca*)? Aquaculture
565 Research, 42(5), 727-732. DOI : 10.1111/j.1365-2109.2010.02681.x

566 Steinfeldt, S., 2015. Culture Methods of Pikeperch Early Life Stages, in: Biology and Culture
567 of Percid Fishes. Springer, pp. 295–312. DOI: 10.1007/978-94-017-7227-3_10

568 Sullivan, J.F., Atchison, G.J., 1978. Predator-prey behavior of fathead minnows, *Pimephales*
569 *promelas* and largemouth bass, *Micropterus salmoides* in a model ecosystem. Journal
570 of Fish Biology 13, 249–253. DOI: 10.1111/j.1095-8649.1978.tb03432.x

571 Szczepkowski, M., Zakęś, Z., Szczepkowska, B., Piotrowska, I., 2011. Effect of size sorting on
572 the survival, growth and cannibalism in pikeperch (*Sander lucioperca* L.) larvae during
573 intensive culture in RAS. Czech Journal of Animal Science, 56(11), 483-489.

574 Takeuchi, N., 2009. Ontogenetic changes in foraging tactics of the piscivorous cornetfish
575 *Fistularia commersonii*. Ichthyological Research, 56(1), 18-27. DOI: 10.1007/s10228-
576 008-0059-9

577 Toomey, L., Bláha, M., Mauduit, E., Vanina, T., Baratçabal, M., Ledoré, Y., ... & Lecocq, T.
578 2019. When behavioural geographic differentiation matters: inter-populational
579 comparison of aggressiveness and group structure in the European perch. Aquaculture
580 International, 27, 1177-1191. DOI: 10.1007/s10499-019-00343-z

581 Toomey, L., Lecocq, T., Bokor, Z., Espinat, L., Ferincz, Á., Goulon, C., ... & Fontaine, P., 2020.
582 Comparison of single-and multi-trait approaches to identify best wild candidates for
583 aquaculture shows that the simple way fails. Scientific Reports, 10(1), 11564. DOI:
584 10.1038/s41598-020-68315-5

585 Turesson, H., Persson, A., Brönmark, C., 2002. Prey size selection in piscivorous pikeperch
586 (*Stizostedion lucioperca*) includes active prey choice. Ecology of Freshwater Fish 11,
587 223–233. DOI: 10.1034/j.1600-0633.2002.00019.x

588 Turesson, H., Brönmark, C., 2004. Foraging behavior and capture success in perch, pikeperch
589 and pike and the effects of prey density. Journal of Fish Biology, 65(2), 363-375. DOI:
590 10.1111/j.0022-1112.2004.00455.x

591 Villacorta, P. J., 2015. ART: Aligned Rank Transform for Nonparametric Factorial Analysis. R
592 package version 1.0. <https://cran.r-project.org/web/packages/ART/index.html>

593 Wang, N., Xu, X.L., Kestemont, P., 2009. Effect of temperature and feeding frequency on
594 growth performances, feed efficiency and body composition of pikeperch juveniles

595 (*Sander lucioperca*). Aquaculture 289 (1-2), 70-73. DOI:
596 10.1016/j.aquaculture.2009.01.002
597 Yang, S., Yang, K., Liu, C., Sun, J., Zhang, F., Zhang, X., Song, Z., 2015. To what extent is
598 cannibalism genetically controlled in fish? A case study in juvenile hybrid catfish *Silurus*
599 *meridionalis-asotus* and the progenitors. Aquaculture, 437, 208-214. DOI:
600 10.1016/j.aquaculture.2014.12.005
601 Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A., Smith, G. M., 2009. Zero-truncated and
602 zero-inflated models for count data, in: Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev,
603 A.A., Smith, G.M. (Eds.), Mixed effects models and extensions in ecology with R.
604 Springer, pp. 261–293. DOI: 10.1007/978-0-387-87458-6_11

605

606 **Funding statement**

607 This work was supported by the European Union’s Seventh Framework Programme for
608 research, technological development and demonstration (KBBE-2013-07 single stage, GA
609 603121, Diversify).

610

611 **Author Contributions**

612 All authors have given their approval to the final version of the manuscript. TC FT PF AP:
613 conceived and designed the experiment. YL: took care of fish and lead routine maintenance
614 of rearing facilities. TC: performed the experiments. TC: analyzed the behavioral video
615 recordings. TC AP: analyzed the behavioral and morphological data. TC YL FT PF AP: wrote
616 the paper.

617

618 **Conflict of interest statement**

619 The authors declare no competing interests.

620

621 **Figures captions**

622

623 **Figure 1:** Comparisons of the number of orientations (A), approaches (B), attacks (C),
624 captures (D), pursuits (E), and performance (F) between larval pikeperch *Sander lucioperca*
625 cannibals and non-cannibals confronted to three pikeperch larvae in cannibalism tests and
626 to three zebrafish *Danio rerio* larvae in predation tests (see definitions in Table 1). The trial
627 duration was 20 minutes after 30 minutes of acclimatization period.

628 Black lines represented median, black triangles mean and black points outliers. Different
629 letters indicated a statistical difference at $p_{\text{adjust}} < 0.05$ between cannibals and non-cannibals
630 in each test (predation and cannibalism) for each behavior.


631

632 **Figure 2:** Comparisons for larval pikeperch *Sander lucioperca* (A) cannibals and (B) non-
633 cannibals between the number of attacks with “S-shape” and without “S-shape” displayed in
634 each test (cannibalism with three pikeperch larvae and predation with three zebrafish *Danio*
635 *rerio* larvae). The trial duration was 20 minutes after 30 minutes of acclimatization period.

636 Black lines represented median, black triangles mean and black points outliers. Different
637 letters indicated a statistical difference at $p_{\text{adjust}} < 0.05$ between the two types of attacks
638 (with or without “S-shape”) in each test (predation and cannibalism) for each status
639 (cannibals and non-cannibals).


640

641 Figure 1.



642
643

644 Figure 2.



645

646

647 **Table 1.** Definitions of studied behaviors in pikeperch larvae.

Behaviors	Definitions
Orientation	Predator turn its head toward prey or conspecific and eye tracked it (Bell and Sih, 2007).
Approach	Predator movement toward prey or conspecific with slow swimming (Colchen et al., 2017).
Attack without “S-shape”	Predator move toward prey or conspecific and attack with a small and fast tail beat, projecting it rapidly (Sullivan and Atchinson, 1978).
Attack with “S-shape”	A fast movement of predator toward prey or conspecific, with an open mouth (Colchen et al., 2019). The attack is marked by a stop of the fish and the formation of a S-shape with his body before a fast movement (Houde, 2001).
Capture	Predator bite to catch the prey or the conspecific (Colchen et al., 2020a).
Pursuit	When capture failed, predator rapidly swim to catch up the prey or the conspecific.

648
649

650 **Table 2.** Impact of the order and status*order on behaviors (number of orientations,
 651 approaches, attacks and captures). The order corresponds to the order in which pikeperch
 652 larvae were tested first cannibalism or predation) and status designs whether the individuals
 653 are either cannibals or non-cannibals.

654

	Orientations	Approaches	Attacks	Captures
Order	$\chi^2 = 1.83;$ p = 0.18	$\chi^2 = 2.14;$ p = 0.14	$\chi^2 = 0.29;$ p = 0.59	$\chi^2 = 0.004;$ p = 0.95
Status x Order	$\chi^2 = 0.99;$ p = 0.32	$\chi^2 = 0.98;$ p = 0.32	$\chi^2 = 0.42;$ p = 0.52	$\chi^2 = 1.25;$ p = 0.26

655

656

657 **Table 3.** Comparisons between predation and cannibalism tests for cannibals and non-
658 cannibals for all behaviors. The trial duration was 20 minutes after 30 minutes of
659 acclimatization period. Each behavior implied in a predation sequence (orientation,
660 approach, attack, capture, pursuit) was represented and the performance was calculated as
661 the number of succeed captures on the total number of attacks. W is the value of Wilcoxon
662 test and the p-value is adjusted. All results were represented by their means and Standard
663 Deviation (SD).
664

Behaviors	Orientation	Approaches	Attacks	Captures	Pursuits	Performance
<i>Cannibals</i>						
Mean ± SD in predation test	14 ± 11	9 ± 7	9 ± 8	2 ± 1	2 ± 3	0.31 ± 0.28
Mean ± SD in cannibalism test	40 ± 28	23 ± 21	4 ± 4	1 ± 1	0 ± 0	0.21 ± 0.29
Comparison between both tests	W = 1185; p_{adjust} < 0.0001	W = 1056; p_{adjust} < 0.0001	W = 148; p_{adjust} < 0.0001	W = 50; p_{adjust} < 0.0001	W = 66.5; p_{adjust} = 0.001	W = 122; p_{adjust} = 0.02
<i>Non-cannibals</i>						
Mean ± SD in predation test	30 ± 20	17 ± 12	19 ± 15	0 ± 1	4 ± 7	0.05 ± 0.15
Mean ± SD in cannibalism test	30 ± 26	17 ± 21	3 ± 6	0 ± 0	0 ± 1	0.02 ± 0.08
Comparison between both tests	W = 517; p_{adjust} = 1	W = 458; p_{adjust} = 0.7	W = 521; p_{adjust} < 0.0001	W = 14; p_{adjust} = 0.004	W = 3.5; p_{adjust} < 0.0001	W = 23; p_{adjust} = 0.4

665
666
667